



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 06:10 PM UTC

PDB ID : 6CMP / pdb_00006cmp
Title : Closed structure of inactive SHP2 mutant C459E
Authors : Padua, R.A.P.; Sun, Y.; Marko, I.; Pitsawong, W.; Kern, D.
Deposited on : 2018-03-06
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	532	<div> <div>7%</div> <div> <div></div> <div>88%</div> <div>6%</div> <div>6%</div> </div> </div>
1	B	532	<div> <div>5%</div> <div> <div></div> <div>86%</div> <div>6%</div> <div>8%</div> </div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 16211 atoms, of which 7559 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosine-protein phosphatase non-receptor type 11.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	498	Total	C	H	N	O	S	0	14	0
			7845	2516	3845	718	748	18			
1	B	489	Total	C	H	N	O	S	0	4	0
			7603	2451	3714	689	732	17			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	GLY	-	expression tag	UNP Q06124
A	-1	SER	-	expression tag	UNP Q06124
A	0	GLY	-	expression tag	UNP Q06124
A	459	GLU	CYS	engineered mutation	UNP Q06124
B	-2	GLY	-	expression tag	UNP Q06124
B	-1	SER	-	expression tag	UNP Q06124
B	0	GLY	-	expression tag	UNP Q06124
B	459	GLU	CYS	engineered mutation	UNP Q06124


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	372	Total	O	0	0
			372	372		
2	B	391	Total	O	0	0
			391	391		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



- Molecule 1: Tyrosine-protein phosphatase non-receptor type 11



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	54.75Å 84.91Å 211.92Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	53.00 – 1.80 53.00 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.2 (53.00-1.80) 89.2 (53.00-1.80)	Depositor EDS
R_{merge}	0.16	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.44 (at 1.81Å)	Xtrriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.201 , 0.231 0.202 , 0.231	Depositor DCC
R_{free} test set	1996 reflections (2.17%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtrriage
Anisotropy	0.395	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 37.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	16211	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.29 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.1108e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/4144	0.38	0/5602
1	B	0.17	0/3992	0.37	0/5407
All	All	0.18	0/8136	0.38	0/11009

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4000	3845	3790	19	0
1	B	3889	3714	3693	15	0
2	A	372	0	0	1	0
2	B	391	0	0	0	0
All	All	8652	7559	7483	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:476:ILE:HD12	1:B:521:TYR:CG	2.37	0.59

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:484:VAL:HG23	1:B:522:ILE:HG23	1.86	0.57
1:A:107:PRO:HG3	1:A:190:LEU:HD12	1.87	0.55
1:B:125:LEU:HB3	1:B:216:LEU:HD21	1.88	0.55
1:A:126:LEU:HD23	1:A:216:LEU:HG	1.90	0.53
1:B:230:VAL:HG13	1:B:516:MET:HE1	1.94	0.50
1:A:125:LEU:HB3	1:A:216:LEU:HD11	1.94	0.48
1:B:355:MET:HE2	1:B:357:THR:HG23	1.96	0.48
1:A:352:VAL:HG11	1:A:442:VAL:HG13	1.96	0.47
1:B:352:VAL:HG11	1:B:442:VAL:HG13	1.95	0.47
1:B:75:ALA:HB2	1:B:258:GLU:HG3	1.97	0.47
1:A:124:LYS:O	1:A:128:GLU:HG3	2.16	0.46
1:A:355:MET:HE2	1:A:357:THR:HG23	1.97	0.46
1:A:290:VAL:HG11	1:A:344:MET:HG3	1.97	0.46
1:A:355:MET:HE2	1:A:357:THR:CG2	2.46	0.45
1:B:356:THR:HG22	1:B:420:PHE:HB3	1.98	0.44
1:A:356:THR:HG22	1:A:420:PHE:HB3	1.99	0.44
1:B:459:GLU:O	1:B:460:SER:C	2.61	0.43
1:A:467:GLY:HA2	1:A:504:MET:HE3	2.01	0.42
1:A:477:ASP:OD1	1:A:480[A]:ARG:NH1	2.53	0.42
1:B:351:ARG:HG3	1:B:449:ILE:HG21	2.02	0.42
1:B:30:LEU:HD12	1:B:30:LEU:C	2.45	0.41
1:A:330:THR:HG23	1:A:458:HIS:HB3	2.02	0.41
1:A:77:LEU:C	1:A:77:LEU:HD23	2.45	0.41
1:A:377:LEU:HD11	1:A:384:ARG:HG2	2.02	0.41
1:B:418:TYR:HB3	1:B:438:PHE:CE1	2.56	0.41
1:A:421:ARG:NH2	2:A:638:HOH:O	2.53	0.41
1:A:459:GLU:O	1:A:460:SER:C	2.64	0.41
1:B:77:LEU:C	1:B:77:LEU:HD23	2.46	0.40
1:B:467:GLY:HA2	1:B:504:MET:HE3	2.02	0.40
1:A:30:LEU:HD12	1:A:30:LEU:C	2.46	0.40
1:A:418:TYR:HB3	1:A:438:PHE:CE1	2.56	0.40
1:B:330:THR:HG23	1:B:458:HIS:HB3	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries

of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	502/532 (94%)	487 (97%)	14 (3%)	1 (0%)	43	31
1	B	481/532 (90%)	468 (97%)	12 (2%)	1 (0%)	43	31
All	All	983/1064 (92%)	955 (97%)	26 (3%)	2 (0%)	43	31

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	505	VAL
1	B	505	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	423/472 (90%)	422 (100%)	1 (0%)	87	87
1	B	414/472 (88%)	412 (100%)	2 (0%)	81	80
All	All	837/944 (89%)	834 (100%)	3 (0%)	84	83

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	216	LEU
1	B	151	VAL
1	B	523	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	57	GLN
1	A	175	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	196	HIS
1	A	426	HIS
1	A	444	HIS
1	A	446	GLN
1	A	506	GLN
1	B	58	ASN
1	B	85	HIS
1	B	200	ASN
1	B	335	GLN
1	B	426	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	498/532 (93%)	0.37	35 (7%)	22 21	13, 33, 62, 86	7 (1%)
1	B	489/532 (91%)	0.33	25 (5%)	33 32	13, 33, 56, 77	2 (0%)
All	All	987/1064 (92%)	0.35	60 (6%)	27 25	13, 33, 58, 86	9 (0%)

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	316	THR	5.6
1	B	92	ASN	4.9
1	A	85	HIS	4.8
1	A	301	VAL	4.4
1	B	142	SER	4.4
1	A	86	GLY	4.0
1	A	142	SER	3.7
1	A	237	ALA	3.7
1	B	143	HIS	3.7
1	A	236	LEU	3.5
1	A	162	ASP	3.5
1	A	90	GLU	3.5
1	B	450	MET	3.4
1	A	160	SER	3.4
1	B	248	TRP	3.3
1	A	158	GLY	3.2
1	A	366	LYS	3.1
1	B	205	THR	3.0
1	A	39	GLY	3.0
1	B	527	ARG	2.9
1	B	86	GLY	2.9
1	A	159	GLU	2.9
1	A	143	HIS	2.8
1	B	525	LEU	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	248	TRP	2.8
1	A	144	PRO	2.8
1	B	314	PHE	2.7
1	A	524	THR	2.7
1	A	450	MET	2.7
1	A	527	ARG	2.7
1	A	481[A]	GLU	2.7
1	B	91	LYS	2.7
1	A	409	GLY	2.6
1	B	89	LYS	2.6
1	A	315	GLU	2.6
1	A	38	PRO	2.6
1	A	155	ASP	2.6
1	A	163	GLY	2.5
1	A	205	THR	2.5
1	B	324	LYS	2.5
1	A	145	GLY	2.4
1	A	84	HIS	2.4
1	B	235	LYS	2.4
1	B	197	TYR	2.3
1	B	357	THR	2.3
1	B	484	VAL	2.3
1	B	3	SER	2.3
1	B	301	VAL	2.3
1	B	144	PRO	2.2
1	A	161	ASN	2.2
1	B	171	MET	2.2
1	B	275	ASN	2.2
1	A	117	LEU	2.2
1	A	484	VAL	2.1
1	A	95	VAL	2.1
1	A	156	ASP	2.1
1	B	409	GLY	2.1
1	B	85	HIS	2.0
1	B	140	SER	2.0
1	A	93	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.