



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 04:29 AM EDT

PDB ID : 9CPU / pdb\_00009cpu  
Title : Crystal structure of SARS-CoV-2 receptor binding domain in complex with antibodies C03-0304 and CC12.3  
Authors : Feng, Z.; Wilson, I.A.  
Deposited on : 2024-07-18  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	4008 (2.60-2.60)
Clashscore	190562	4347 (2.60-2.60)
Ramachandran outliers	187476	4277 (2.60-2.60)
Sidechain outliers	187428	4277 (2.60-2.60)
RSRZ outliers	180081	4008 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	 4% 86% 9% .
2	H	222	 3% 90% 8% .
3	F	220	 11% 85% 11% .
4	L	220	 % 92% 8% .
5	G	214	 3% 82% 16% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
6	B	3	 67% 33%

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 8521 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Spike protein S1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1552	995	259	290	8			

There are 7 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	531	GLY	-	expression tag	UNP P0DTC2
A	532	HIS	-	expression tag	UNP P0DTC2
A	533	HIS	-	expression tag	UNP P0DTC2
A	534	HIS	-	expression tag	UNP P0DTC2
A	535	HIS	-	expression tag	UNP P0DTC2
A	536	HIS	-	expression tag	UNP P0DTC2
A	537	HIS	-	expression tag	UNP P0DTC2

- Molecule 2 is a protein called C03-0304 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	H	216	Total	C	N	O	S	0	0	0
			1605	1018	261	318	8			

- Molecule 3 is a protein called CC12.3 Fab heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	F	212	Total	C	N	O	S	0	0	0
			1596	1014	263	313	6			


- Molecule 4 is a protein called C03-0304 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	L	219	Total	C	N	O	S	0	0	0
			1703	1071	283	344	5			

- Molecule 5 is a protein called CC12.3 Fab light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	G	211	Total	C	N	O	S	0	0	0
			1622	1014	277	327	4			

- Molecule 6 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	B	3	Total	C	N	O		0	0	0
			39	22	2	15				


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	83	Total	O	0	0
			83	83		
7	H	93	Total	O	0	0
			93	93		
7	F	72	Total	O	0	0
			72	72		
7	L	101	Total	O	0	0
			101	101		
7	G	55	Total	O	0	0
			55	55		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Spike protein S1




- Molecule 2: C03-0304 Fab heavy chain




- Molecule 3: CC12.3 Fab heavy chain




- Molecule 4: C03-0304 Fab light chain



- Molecule 5: CC12.3 Fab light chain





● Molecule 6: beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	110.23Å 110.23Å 230.35Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.30 – 2.60 49.30 – 2.60	Depositor EDS
% Data completeness (in resolution range)	98.8 (49.30-2.60) 98.9 (49.30-2.60)	Depositor EDS
$R_{merge}$	0.10	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 2.58Å)	Xtriage
Refinement program	PHENIX (1.21rc1_5127: ???)	Depositor
R, $R_{free}$	0.206 , 0.239 0.207 , 0.239	Depositor DCC
$R_{free}$ test set	2000 reflections (4.47%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	37.1	Xtriage
Anisotropy	0.047	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.28 , 31.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	8521	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.63% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.07	0/1596	0.24	0/2172
2	H	0.08	0/1646	0.27	0/2243
3	F	0.10	0/1635	0.27	0/2226
4	L	0.08	0/1741	0.26	0/2364
5	G	0.09	0/1657	0.29	0/2249
All	All	0.09	0/8275	0.27	0/11254

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1552	0	1472	12	0
2	H	1605	0	1575	11	0
3	F	1596	0	1557	14	0
4	L	1703	0	1652	8	0
5	G	1622	0	1583	26	0
6	B	39	0	34	0	0
7	A	83	0	0	0	0
7	F	72	0	0	1	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
7	G	55	0	0	8	0
7	H	93	0	0	2	0
7	L	101	0	0	1	0
All	All	8521	0	7873	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:G:142:ARG:NH2	7:G:304:HOH:O	2.23	0.70
2:H:126:PRO:HG3	2:H:138:LEU:HB3	1.72	0.69
5:G:76:SER:O	7:G:301:HOH:O	2.12	0.67
5:G:124:GLN:O	7:G:302:HOH:O	2.13	0.67
2:H:81:GLN:OE1	7:H:301:HOH:O	2.16	0.62
5:G:190:LYS:HE3	5:G:210:ASN:HD22	1.63	0.62
5:G:90:GLN:OE1	7:G:303:HOH:O	2.16	0.60
2:H:163:VAL:HG22	2:H:182:VAL:HG22	1.85	0.58
5:G:24:ARG:HH22	5:G:69:THR:HG23	1.70	0.56
5:G:21:LEU:HD12	5:G:73:LEU:HD23	1.88	0.56
5:G:145:LYS:HB3	5:G:197:THR:HB	1.87	0.56
2:H:97:GLY:O	2:H:100:THR:OG1	2.23	0.55
2:H:159:LEU:HD21	2:H:182:VAL:HG11	1.89	0.54
5:G:128:GLY:N	7:G:302:HOH:O	2.41	0.54
2:H:70:SER:OG	7:H:301:HOH:O	2.18	0.54
4:L:120:PRO:HD3	4:L:132:VAL:HG22	1.90	0.53
5:G:129:THR:N	7:G:302:HOH:O	2.32	0.53
1:A:360:ASN:HA	1:A:523:THR:HB	1.91	0.53
4:L:37:GLN:HB2	4:L:47:LEU:HD11	1.91	0.52
1:A:520:ALA:HB1	1:A:521:PRO:HD2	1.91	0.52
1:A:334:ASN:N	1:A:334:ASN:OD1	2.43	0.51
3:F:119:PRO:HB3	3:F:145:TYR:HB3	1.93	0.51
5:G:54:ARG:NH1	7:G:309:HOH:O	2.44	0.50
3:F:82:MET:HE1	3:F:109:VAL:HG21	1.94	0.50
4:L:142:ARG:HD3	4:L:163:VAL:HG21	1.92	0.50
1:A:389:ASP:N	1:A:389:ASP:OD1	2.42	0.50
1:A:379:CYS:SG	1:A:384:PRO:HG3	2.51	0.50
3:F:184:VAL:HG22	3:F:185:PRO:HD2	1.94	0.50
1:A:409:GLN:HA	1:A:414:GLN:HG2	1.95	0.49
4:L:80:ALA:HA	4:L:106:ILE:HD11	1.94	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:L:186:TYR:O	4:L:192:TYR:OH	2.29	0.48
5:G:37:GLN:HB2	5:G:47:LEU:HD11	1.96	0.47
5:G:170:ASP:OD1	5:G:170:ASP:N	2.46	0.47
5:G:59:PRO:HB2	5:G:61:ARG:HG2	1.97	0.46
5:G:24:ARG:NH2	5:G:69:THR:HG23	2.30	0.46
3:F:72:ASP:HB2	3:F:79:TYR:HE2	1.80	0.46
3:F:39:GLN:HB2	3:F:45:LEU:HD23	1.97	0.46
4:L:210:ASN:HB2	4:L:213:GLU:HB2	1.97	0.45
5:G:93:SER:N	7:G:303:HOH:O	2.38	0.45
3:F:11:LEU:HB2	3:F:147:PRO:HG3	1.98	0.44
5:G:183:LYS:HB3	5:G:183:LYS:HE3	1.73	0.44
5:G:190:LYS:HE3	5:G:210:ASN:ND2	2.30	0.44
2:H:93:ALA:HB1	2:H:100(B):MET:HB3	1.99	0.44
3:F:87:THR:HG23	3:F:110:THR:HA	2.00	0.44
4:L:183:LYS:O	4:L:187:GLU:HG3	2.17	0.44
5:G:185:ASP:HA	5:G:188:LYS:HD2	2.00	0.44
4:L:18:ARG:NH1	7:L:315:HOH:O	2.50	0.44
5:G:14:SER:O	5:G:17:GLU:HG3	2.17	0.44
1:A:377:PHE:CD1	2:H:31:THR:HG22	2.53	0.43
1:A:502:GLY:O	1:A:506:GLN:HG3	2.18	0.43
2:H:43:LYS:HA	2:H:43:LYS:HD3	1.90	0.43
2:H:119:PRO:HB3	2:H:145:TYR:HB3	2.01	0.43
3:F:143:LYS:NZ	7:F:307:HOH:O	2.44	0.43
1:A:489:TYR:OH	3:F:94:ARG:NH2	2.53	0.42
3:F:154:TRP:CZ3	3:F:196:CYS:HB3	2.55	0.42
5:G:24:ARG:HE	5:G:70:ASP:CG	2.28	0.42
3:F:52:TYR:O	3:F:71:ARG:NH2	2.44	0.42
5:G:148:TRP:CD2	5:G:179:LEU:HD12	2.55	0.42
1:A:415:THR:HG22	3:F:58:PHE:CZ	2.55	0.41
1:A:390:LEU:HD23	1:A:390:LEU:HA	1.80	0.41
5:G:80:PRO:HA	5:G:106:ILE:HG13	2.03	0.41
1:A:376:THR:HG23	1:A:378:LYS:HG3	2.02	0.41
5:G:54:ARG:HD3	5:G:62:PHE:O	2.20	0.41
3:F:33:TYR:CE1	3:F:97:GLY:HA2	2.56	0.41
2:H:35:GLY:HA2	2:H:50:ILE:HA	2.02	0.41
5:G:207:LYS:HD3	5:G:207:LYS:HA	1.93	0.41
3:F:42:GLY:C	3:F:43:LYS:HD2	2.46	0.40
5:G:36:TYR:HE1	5:G:89:GLN:HB3	1.87	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/205 (95%)	191 (98%)	3 (2%)	0	100	100
2	H	212/222 (96%)	210 (99%)	2 (1%)	0	100	100
3	F	208/220 (94%)	202 (97%)	6 (3%)	0	100	100
4	L	217/220 (99%)	211 (97%)	6 (3%)	0	100	100
5	G	209/214 (98%)	203 (97%)	6 (3%)	0	100	100
All	All	1040/1081 (96%)	1017 (98%)	23 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	169/177 (96%)	168 (99%)	1 (1%)	78	91
2	H	182/188 (97%)	182 (100%)	0	100	100
3	F	180/186 (97%)	178 (99%)	2 (1%)	65	84
4	L	193/194 (100%)	191 (99%)	2 (1%)	68	86
5	G	183/185 (99%)	182 (100%)	1 (0%)	81	92
All	All	907/930 (98%)	901 (99%)	6 (1%)	76	89

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	494	SER
3	F	184	VAL
3	F	189	LEU
4	L	55	GLU
4	L	78	LEU
5	G	70	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
2	H	81	GLN
3	F	3	GLN
3	F	164	HIS
3	F	199	ASN
4	L	152	ASN
4	L	160	GLN
5	G	137	ASN
5	G	189	HIS

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
6	NAG	B	1	1,6	14,14,15	0.74	0	17,19,21	0.93	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
6	NAG	B	2	6	14,14,15	0.70	0	17,19,21	0.86	0
6	BMA	B	3	6	11,11,12	0.82	0	15,15,17	1.77	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	B	1	1,6	-	0/6/23/26	0/1/1/1
6	NAG	B	2	6	-	0/6/23/26	0/1/1/1
6	BMA	B	3	6	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	B	3	BMA	C1-O5-C5	5.61	119.71	112.19


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	196/205 (95%)	0.24	9 (4%) 37 32	16, 34, 67, 121	0
2	H	216/222 (97%)	-0.04	7 (3%) 50 44	18, 33, 70, 97	0
3	F	212/220 (96%)	0.49	24 (11%) 10 8	20, 40, 90, 109	0
4	L	219/220 (99%)	0.04	3 (1%) 73 69	19, 35, 56, 75	0
5	G	211/214 (98%)	0.63	6 (2%) 55 49	25, 48, 70, 113	0
All	All	1054/1081 (97%)	0.27	49 (4%) 37 32	16, 37, 77, 121	0

All (49) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
5	G	169	LYS	4.2
3	F	189	LEU	3.9
5	G	168	SER	3.4
3	F	190	GLY	3.4
3	F	159	LEU	3.4
5	G	23	CYS	3.4
2	H	133	GLY	3.2
3	F	187	SER	3.0
3	F	214	LYS	2.9
2	H	132	SER	2.8
1	A	517	LEU	2.8
3	F	128	SER	2.7
1	A	528	LYS	2.7
3	F	135	THR	2.7
1	A	521	PRO	2.7
1	A	520	ALA	2.7
3	F	186	SER	2.6
4	L	62	PHE	2.6
1	A	518	LEU	2.6
3	F	156	SER	2.6

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	346	ARG	2.6
2	H	213	PRO	2.6
3	F	185	PRO	2.6
5	G	51	ALA	2.5
3	F	140	CYS	2.5
4	L	27(F)	SER	2.5
3	F	211	VAL	2.4
3	F	183	THR	2.4
3	F	196	CYS	2.4
3	F	212	GLU	2.4
5	G	1	GLU	2.4
1	A	519	HIS	2.3
3	F	157	GLY	2.3
1	A	334	ASN	2.3
3	F	158	ALA	2.3
3	F	188	SER	2.3
3	F	184	VAL	2.3
2	H	190	GLY	2.2
1	A	522	ALA	2.2
2	H	185	PRO	2.2
2	H	189	LEU	2.2
4	L	213	GLU	2.1
3	F	95	ASP	2.1
3	F	162	GLY	2.1
2	H	128	SER	2.1
3	F	125	ALA	2.1
3	F	194	TYR	2.1
3	F	161	SER	2.0
5	G	106	ILE	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
6	NAG	B	1	14/15	-	-	29,35,44,44	0
6	NAG	B	2	14/15	-	-	36,53,64,76	0
6	BMA	B	3	11/12	-	-	65,75,85,89	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.