



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 1, 2026 – 08:07 AM UTC

PDB ID : 4E79 / pdb_00004e79
Title : Structure of LpxD from *Acinetobacter baumannii* at 2.66Å resolution (P4322 form)
Authors : Badger, J.; Chie-Leon, B.; Logan, C.; Sridhar, V.; Sankaran, B.; Zwart, P.H.; Nienaber, V.
Deposited on : 2012-03-16
Resolution : 2.66 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.66 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1110 (2.66-2.66)
Clashscore	190562	1141 (2.66-2.66)
Ramachandran outliers	187476	1126 (2.66-2.66)
Sidechain outliers	187428	1126 (2.66-2.66)
RSRZ outliers	180081	1110 (2.66-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	357	 75% 19% ...
1	B	357	 76% 20% ..
1	C	357	 80% 15% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7722 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called UDP-3-O-acylglucosamine N-acyltransferase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	346	Total	C	N	O	S	0	0	0
			2560	1602	451	499	8			
1	B	350	Total	C	N	O	S	0	0	0
			2597	1623	458	508	8			
1	C	348	Total	C	N	O	S	0	0	0
			2552	1599	450	495	8			

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	GLY	-	expression tag	UNP B0VMV2
A	2	GLY	-	expression tag	UNP B0VMV2
A	3	SER	-	expression tag	UNP B0VMV2
B	1	GLY	-	expression tag	UNP B0VMV2
B	2	GLY	-	expression tag	UNP B0VMV2
B	3	SER	-	expression tag	UNP B0VMV2
C	1	GLY	-	expression tag	UNP B0VMV2
C	2	GLY	-	expression tag	UNP B0VMV2
C	3	SER	-	expression tag	UNP B0VMV2


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	6	Total	O	0	0
			6	6		
2	B	4	Total	O	0	0
			4	4		
2	C	3	Total	O	0	0
			3	3		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: UDP-3-O-acylglucosamine N-acyltransferase



• Molecule 1: UDP-3-O-acylglucosamine N-acyltransferase



• Molecule 1: UDP-3-O-acylglucosamine N-acyltransferase



Q344				
	L350			
	E351			
	S352			
	THR			
	PHE			
	ASN			
	LEU			
	ARG			

4 Data and refinement statistics

Property	Value	Source
Space group	P 43 2 2	Depositor
Cell constants a, b, c, α , β , γ	64.67Å 64.67Å 463.81Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.60 – 2.66 49.60 – 2.66	Depositor EDS
% Data completeness (in resolution range)	(Not available) (49.60-2.66) 99.4 (49.60-2.66)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.17 (at 2.65Å)	Xtriage
Refinement program	REFMAC 5.5.0072	Depositor
R, R_{free}	0.234 , 0.300 0.227 , 0.292	Depositor DCC
R_{free} test set	1508 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	43.9	Xtriage
Anisotropy	0.362	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 20.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	7722	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.68	0/2600	0.97	3/3539 (0.1%)
1	B	0.66	0/2637	0.92	0/3587
1	C	0.69	0/2592	0.94	2/3529 (0.1%)
All	All	0.68	0/7829	0.94	5/10655 (0.0%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	179	SER	N-CA-C	5.79	118.41	110.24
1	C	311	GLY	N-CA-C	-5.34	106.31	112.29
1	A	146	LYS	O-C-N	5.31	129.75	123.27
1	A	251	ASN	N-CA-C	5.24	119.51	113.38
1	C	38	ALA	N-CA-C	5.11	117.64	110.23

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2560	0	2481	48	0
1	B	2597	0	2529	50	0
1	C	2552	0	2460	39	0
2	A	6	0	0	0	0
2	B	4	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	3	0	0	0	0
All	All	7722	0	7470	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:39:GLU:H	1:B:42:HIS:HD2	1.06	0.95
1:B:39:GLU:H	1:B:42:HIS:CD2	1.94	0.85
1:C:150:ASN:HB2	1:C:168:SER:HB3	1.63	0.80
1:B:333:LEU:HD13	1:B:334:THR:H	1.51	0.76
1:A:105:ILE:HG12	1:A:123:ILE:HD12	1.73	0.70
1:A:190:PHE:HZ	1:C:213:ARG:HD3	1.58	0.69
1:A:324:ARG:HB3	1:C:328:LEU:HD13	1.76	0.68
1:C:63:TYR:CE1	1:C:76:LYS:HG2	2.30	0.66
1:B:309:GLY:CA	1:C:295:MET:HE1	2.25	0.66
1:A:190:PHE:CZ	1:C:213:ARG:HD3	2.32	0.64
1:B:35:LEU:HD22	1:B:55:ALA:HA	1.80	0.63
1:C:39:GLU:H	1:C:42:HIS:HD2	1.48	0.61
1:A:66:THR:HG22	1:A:69:LEU:HB2	1.82	0.61
1:A:293:MET:O	1:C:293:MET:HE1	2.01	0.60
1:B:225:LEU:O	1:B:226:ASP:HB2	2.02	0.59
1:B:134:VAL:HB	1:B:152:GLU:HG2	1.84	0.59
1:B:313:PHE:HB2	1:B:318:TRP:HB2	1.83	0.58
1:B:333:LEU:HD13	1:B:334:THR:N	2.18	0.58
1:B:213:ARG:HD3	1:C:190:PHE:HZ	1.69	0.57
1:C:39:GLU:H	1:C:42:HIS:CD2	2.22	0.57
1:B:333:LEU:CD1	1:B:334:THR:H	2.17	0.57
1:B:213:ARG:NH2	1:B:237:ASP:HA	2.21	0.56
1:B:309:GLY:HA3	1:C:295:MET:HE1	1.87	0.56
1:A:39:GLU:H	1:A:42:HIS:HD2	1.52	0.56
1:A:117:ILE:HG12	1:A:123:ILE:HD11	1.88	0.55
1:A:10:LEU:HG	1:A:29:PHE:CE2	2.41	0.55
1:A:314:GLU:CD	1:A:316:ASN:H	2.14	0.55
1:A:225:LEU:HD11	1:C:89:PHE:CZ	2.42	0.55
1:C:150:ASN:CB	1:C:168:SER:HB3	2.37	0.54
1:C:41:ASN:ND2	1:C:42:HIS:CE1	2.75	0.54
1:A:63:TYR:CE1	1:A:76:LYS:HG2	2.41	0.54
1:C:204:GLY:HA3	1:C:226:ASP:O	2.07	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:332:PRO:O	1:B:336:ILE:HG13	2.08	0.54
1:A:39:GLU:H	1:A:42:HIS:CD2	2.26	0.54
1:A:159:ILE:HD13	1:A:165:ILE:HD11	1.89	0.53
1:A:333:LEU:HD22	1:B:324:ARG:HG3	1.91	0.53
1:B:332:PRO:HA	1:B:333:LEU:HD12	1.91	0.53
1:A:9:ARG:HB2	1:A:12:GLU:HG3	1.90	0.52
1:C:315:ASN:O	1:C:318:TRP:HB3	2.09	0.52
1:A:333:LEU:HG	1:B:331:VAL:HG21	1.90	0.52
1:B:10:LEU:HB3	1:B:25:GLY:HA2	1.92	0.51
1:A:146:LYS:HE3	1:C:161:SER:OG	2.10	0.51
1:B:22:ILE:HD12	1:B:81:ILE:HD11	1.93	0.51
1:A:66:THR:HG22	1:A:69:LEU:H	1.76	0.51
1:C:172:ARG:HB3	1:C:210:ASN:OD1	2.12	0.50
1:C:217:ASN:O	1:C:239:LEU:HA	2.10	0.50
1:B:341:ASP:OD2	1:C:339:ARG:NH1	2.45	0.50
1:B:66:THR:HG22	1:B:69:LEU:H	1.76	0.50
1:B:329:ALA:HA	1:C:324:ARG:HD2	1.94	0.50
1:B:332:PRO:C	1:B:333:LEU:HD12	2.37	0.50
1:C:35:LEU:HD22	1:C:55:ALA:HA	1.94	0.50
1:A:152:GLU:HB2	1:A:170:LYS:HG3	1.93	0.49
1:B:13:LEU:HD22	1:B:92:LEU:HD22	1.94	0.49
1:A:328:LEU:HB3	1:B:324:ARG:HG2	1.95	0.49
1:C:66:THR:HG22	1:C:69:LEU:HB2	1.95	0.49
1:A:145:THR:HG23	1:A:163:VAL:O	2.13	0.48
1:B:294:SER:HA	1:B:308:SER:OG	2.14	0.48
1:C:143:SER:O	1:C:144:HIS:HB2	2.13	0.47
1:A:151:VAL:HG21	1:A:165:ILE:O	2.15	0.47
1:B:205:SER:OG	1:B:206:VAL:N	2.47	0.47
1:B:11:ASP:HA	1:B:21:LEU:HD13	1.96	0.47
1:C:7:GLN:HB3	1:C:28:GLN:HB3	1.97	0.46
1:A:30:SER:OG	1:A:41:ASN:ND2	2.48	0.46
1:B:39:GLU:N	1:B:42:HIS:HD2	1.91	0.46
1:B:194:GLN:C	1:B:196:LYS:H	2.24	0.46
1:C:177:ILE:HD13	1:C:183:ILE:HG13	1.98	0.45
1:A:49:GLU:H	1:A:49:GLU:CD	2.24	0.45
1:A:322:ILE:HA	1:A:325:LEU:HD12	1.97	0.45
1:C:167:GLY:O	1:C:205:SER:OG	2.30	0.45
1:A:66:THR:CG2	1:A:69:LEU:H	2.30	0.45
1:B:63:TYR:CZ	1:B:76:LYS:HE3	2.52	0.45
1:B:204:GLY:O	1:B:222:ARG:HD2	2.16	0.45
1:A:10:LEU:HG	1:A:29:PHE:HE2	1.81	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:162:TYR:O	1:A:180:SER:HA	2.17	0.44
1:C:239:LEU:O	1:C:257:LYS:HA	2.16	0.44
1:B:63:TYR:OH	1:B:76:LYS:HE3	2.18	0.44
1:A:242:ILE:HD12	1:A:248:ILE:HD11	1.99	0.44
1:A:319:LYS:O	1:A:323:VAL:HG23	2.17	0.44
1:C:164:THR:HB	1:C:182:VAL:HG13	1.99	0.44
1:C:17:VAL:HG13	1:C:92:LEU:HD21	2.00	0.43
1:A:90:ALA:CB	1:B:200:ILE:HG12	2.47	0.43
1:A:120:THR:O	1:A:138:ASN:HA	2.18	0.43
1:A:271:ILE:O	1:A:289:THR:HA	2.17	0.43
1:B:68:ALA:O	1:B:72:HIS:HD2	2.01	0.43
1:A:171:LEU:HD11	1:A:208:ILE:HD12	1.99	0.43
1:A:292:GLY:HA3	1:B:295:MET:HE2	2.00	0.43
1:A:178:HIS:HB2	1:A:215:GLY:C	2.43	0.43
1:B:63:TYR:CE1	1:B:76:LYS:HG2	2.53	0.42
1:B:112:HIS:CE1	1:B:131:GLU:HG2	2.54	0.42
1:B:279:ALA:HB3	1:B:282:LEU:HD11	2.01	0.42
1:B:213:ARG:HD3	1:C:190:PHE:CZ	2.53	0.42
1:C:167:GLY:O	1:C:168:SER:C	2.62	0.42
1:B:235:ILE:HG13	1:B:253:ALA:HA	2.01	0.42
1:C:16:LEU:HD12	1:C:16:LEU:HA	1.94	0.42
1:A:313:PHE:HB2	1:A:318:TRP:HB2	2.01	0.42
1:A:315:ASN:HB2	1:B:305:THR:HG22	2.01	0.41
1:A:32:LEU:HG	1:A:96:PHE:CD1	2.55	0.41
1:A:237:ASP:HB3	1:A:256:ALA:N	2.35	0.41
1:A:255:ALA:HB3	1:A:273:ALA:C	2.46	0.41
1:B:193:TYR:CE1	1:B:198:HIS:HB2	2.56	0.41
1:A:169:SER:OG	1:A:206:VAL:HB	2.20	0.41
1:B:242:ILE:HD12	1:B:248:ILE:HD11	2.03	0.41
1:B:111:ILE:HG21	1:B:117:ILE:HD11	2.03	0.41
1:B:213:ARG:HH21	1:B:237:ASP:HA	1.86	0.41
1:B:309:GLY:HA2	1:C:295:MET:HE1	2.00	0.41
1:C:275:ALA:O	1:C:293:MET:HA	2.21	0.41
1:A:57:VAL:O	1:A:57:VAL:CG1	2.69	0.40
1:A:225:LEU:CD1	1:C:89:PHE:CZ	3.04	0.40
1:A:324:ARG:NH1	1:C:329:ALA:O	2.55	0.40
1:A:343:ILE:HD12	1:C:340:LEU:HD22	2.02	0.40
1:B:328:LEU:HD22	1:C:324:ARG:HG2	2.03	0.40
1:B:52:LEU:HD22	1:B:56:LYS:HE2	2.03	0.40
1:A:34:SER:O	1:A:38:ALA:HB2	2.21	0.40
1:B:332:PRO:HD2	1:B:336:ILE:HD11	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	344/357 (96%)	323 (94%)	19 (6%)	2 (1%)	21	34
1	B	348/357 (98%)	335 (96%)	11 (3%)	2 (1%)	21	34
1	C	346/357 (97%)	333 (96%)	11 (3%)	2 (1%)	21	34
All	All	1038/1071 (97%)	991 (96%)	41 (4%)	6 (1%)	21	34

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	333	LEU
1	C	172	ARG
1	C	250	SER
1	A	225	LEU
1	B	172	ARG
1	A	154	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	268/295 (91%)	249 (93%)	19 (7%)	13	23
1	B	275/295 (93%)	251 (91%)	24 (9%)	9	16
1	C	263/295 (89%)	241 (92%)	22 (8%)	10	17

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
All	All	806/885 (91%)	741 (92%)	65 (8%)	11	18

All (65) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	VAL
1	A	10	LEU
1	A	16	LEU
1	A	30	SER
1	A	40	VAL
1	A	59	ARG
1	A	66	THR
1	A	73	LEU
1	A	76	LYS
1	A	117	ILE
1	A	146	LYS
1	A	165	ILE
1	A	168	SER
1	A	172	ARG
1	A	225	LEU
1	A	272	LEU
1	A	319	LYS
1	A	334	THR
1	A	348	GLU
1	B	10	LEU
1	B	16	LEU
1	B	21	LEU
1	B	27	LEU
1	B	40	VAL
1	B	52	LEU
1	B	66	THR
1	B	103	THR
1	B	123	ILE
1	B	139	THR
1	B	177	ILE
1	B	194	GLN
1	B	203	LEU
1	B	207	LEU
1	B	265	LYS
1	B	293	MET
1	B	298	LYS
1	B	324	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	326	ARG
1	B	328	LEU
1	B	333	LEU
1	B	337	THR
1	B	339	ARG
1	B	350	LEU
1	C	10	LEU
1	C	16	LEU
1	C	30	SER
1	C	40	VAL
1	C	56	LYS
1	C	57	VAL
1	C	66	THR
1	C	69	LEU
1	C	73	LEU
1	C	76	LYS
1	C	139	THR
1	C	168	SER
1	C	181	THR
1	C	207	LEU
1	C	265	LYS
1	C	324	ARG
1	C	325	LEU
1	C	328	LEU
1	C	334	THR
1	C	337	THR
1	C	343	ILE
1	C	344	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (20) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	ASN
1	A	42	HIS
1	A	287	ASN
1	A	346	GLN
1	B	28	GLN
1	B	37	ASN
1	B	42	HIS
1	B	72	HIS
1	B	138	ASN
1	B	202	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	227	ASN
1	B	287	ASN
1	B	344	GLN
1	C	15	HIS
1	C	28	GLN
1	C	41	ASN
1	C	42	HIS
1	C	54	GLN
1	C	144	HIS
1	C	344	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	346/357 (96%)	0.04	4 (1%) 76 72	26, 41, 60, 88	0
1	B	350/357 (98%)	0.03	4 (1%) 78 75	29, 42, 60, 89	0
1	C	348/357 (97%)	0.10	8 (2%) 61 54	27, 41, 69, 90	0
All	All	1044/1071 (97%)	0.06	16 (1%) 72 67	26, 42, 61, 90	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	351	GLU	4.1
1	C	352	SER	3.9
1	C	47	ASN	3.7
1	A	5	VAL	3.1
1	C	350	LEU	3.1
1	B	350	LEU	2.6
1	C	341	ASP	2.6
1	A	335	GLN	2.5
1	B	195	GLY	2.4
1	B	89	PHE	2.3
1	B	354	PHE	2.3
1	C	333	LEU	2.3
1	C	193	TYR	2.2
1	A	349	SER	2.2
1	A	350	LEU	2.1
1	C	338	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.