



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:43 AM UTC

PDB ID : 7ECT / pdb_00007ect
Title : Crystal Structure of Aspergillus terreus Glutamate Dehydrogenase (AtGDH)
Complexed With Tartrate and NADPH
Authors : Godsora, B.K.J.; Prakash, P.; Puneekar, N.S.; Bhaumik, P.
Deposited on : 2021-03-13
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION




The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	460	
1	B	460	
1	C	460	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	TLA	B	503	-	X	-	-
4	TLA	C	508	-	X	-	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 10712 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glutamate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	459	Total	C	N	O	S	0	0	0
			3459	2183	602	662	12			
1	B	459	Total	C	N	O	S	0	0	0
			3455	2180	601	662	12			
1	C	460	Total	C	N	O	S	0	1	0
			3465	2185	603	664	13			

- Molecule 2 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃) (labeled as "Ligand of Interest" by depositor).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	C	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	B	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is L(+)-TARTARIC ACID (CCD ID: TLA) (formula: $C_4H_6O_6$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	4	6		
4	B	1	Total	C	O	0	0
			10	4	6		
4	C	1	Total	C	O	0	0
			10	4	6		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	19	Total	O	0	0
			19	19		
5	B	40	Total	O	0	0
			40	40		
5	C	28	Total	O	0	0
			28	28		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Glutamate dehydrogenase



- Molecule 1: Glutamate dehydrogenase



- Molecule 1: Glutamate dehydrogenase



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	119.86Å 153.09Å 259.27Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.29 – 2.90 39.29 – 2.90	Depositor EDS
% Data completeness (in resolution range)	99.8 (39.29-2.90) 99.8 (39.29-2.90)	Depositor EDS
R_{merge}	0.42	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 2.90Å)	Xtriage
Refinement program	REFMAC 5.8.0189	Depositor
R, R_{free}	0.223 , 0.269 (Not available) , 0.226	Depositor DCC
R_{free} test set	2655 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtriage
Anisotropy	0.179	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 31.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	10712	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NDP, TLA, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.82	0/3529	1.06	3/4774 (0.1%)
1	B	0.82	0/3525	1.04	4/4770 (0.1%)
1	C	0.82	0/3535	1.06	1/4782 (0.0%)
All	All	0.82	0/10589	1.05	8/14326 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	196	ALA	N-CA-C	5.78	117.27	110.97
1	A	185	GLY	N-CA-C	5.58	119.64	112.83
1	C	100	ILE	CB-CA-C	-5.50	104.84	111.88
1	A	196	ALA	N-CA-C	5.37	117.13	111.28
1	B	87	VAL	N-CA-C	5.25	116.02	108.93
1	B	154	ASP	N-CA-C	-5.20	99.72	110.80
1	A	280	VAL	N-CA-C	-5.14	105.39	111.00
1	B	378	ALA	N-CA-C	5.11	116.93	111.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3459	0	3400	23	0
1	B	3455	0	3389	20	0
1	C	3465	0	3396	18	0
2	A	48	0	26	1	0
2	B	48	0	26	0	0
2	C	48	0	26	0	0
3	A	30	0	40	1	0
3	B	6	0	8	0	0
3	C	36	0	48	0	0
4	A	10	0	4	0	0
4	B	10	0	4	0	0
4	C	10	0	4	0	0
5	A	19	0	0	0	0
5	B	40	0	0	0	0
5	C	28	0	0	0	0
All	All	10712	0	10371	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:130:ARG:HA	1:C:164:TYR:CD2	2.32	0.64
1:B:311:GLY:O	1:B:336:ALA:HB1	1.99	0.63
1:C:22:GLU:OE1	1:C:34:ARG:NH1	2.28	0.63
1:A:54:GLU:O	1:A:131:ARG:HD2	2.02	0.60
1:B:223:ARG:NH1	1:B:312:ALA:HB1	2.18	0.59
1:A:82:ARG:HH12	1:A:322:GLN:HE22	1.49	0.58
1:B:224:VAL:HG13	1:B:315:VAL:HB	1.89	0.54
1:B:193:ARG:O	1:B:194:PRO:C	2.54	0.51
1:B:55:ASN:HD21	1:B:59:GLU:HB2	1.74	0.50
1:C:192:ILE:HG22	1:C:192:ILE:O	2.12	0.50
1:A:15:LYS:O	1:A:19:SER:HB2	2.12	0.49
1:C:16:GLU:HB2	1:C:351:GLN:HE22	1.78	0.49
1:A:170:ARG:NH2	1:C:72:SER:OG	2.43	0.48
1:B:55:ASN:ND2	1:B:59:GLU:HB2	2.28	0.47
1:C:383:VAL:HA	1:C:386:SER:HB2	1.95	0.47
1:B:203:TYR:CE2	1:B:409:LYS:HG2	2.49	0.47
1:B:163:GLY:HA2	1:B:187:TRP:CH2	2.49	0.47
1:C:108:LEU:HB2	1:C:110:MET:HE2	1.96	0.47
1:C:192:ILE:O	1:C:196:ALA:HB2	2.15	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:224:VAL:HG21	1:B:240:VAL:HG11	1.98	0.46
1:A:226:ILE:HD12	1:A:247:VAL:HG13	1.97	0.46
1:A:227:SER:OG	1:A:318:PRO:HA	2.16	0.45
1:C:330:ALA:HB1	1:C:357:PHE:CE1	2.52	0.45
1:B:341:ILE:HD13	1:B:357:PHE:CE2	2.52	0.45
1:B:341:ILE:HD13	1:B:357:PHE:CZ	2.52	0.45
1:B:169:TYR:C	1:B:169:TYR:CD1	2.94	0.44
1:B:227:SER:OG	1:B:318:PRO:HA	2.18	0.44
1:A:94:PHE:CD1	1:A:94:PHE:C	2.95	0.44
1:B:383:VAL:HA	1:B:386:SER:HB2	1.99	0.44
1:A:322:GLN:H	1:A:322:GLN:HE21	1.65	0.44
1:A:456:GLN:HA	1:B:163:GLY:O	2.17	0.44
1:C:201:VAL:O	1:C:205:VAL:HG23	2.18	0.44
1:B:141:CYS:SG	1:B:173:ARG:HG2	2.58	0.44
1:C:138:THR:HA	1:C:172:LEU:CD1	2.48	0.43
1:A:195:GLU:HG3	1:A:235:TYR:CE1	2.54	0.43
1:A:195:GLU:HG3	1:A:235:TYR:CD1	2.54	0.43
1:C:72:SER:HB3	1:C:77:TYR:CE1	2.54	0.43
1:A:288:LEU:C	1:A:288:LEU:HD12	2.44	0.43
1:A:361:ARG:NH2	1:A:434:LEU:O	2.51	0.42
2:A:501:NDP:H2A	3:A:504:GOL:C1	2.49	0.42
1:B:339:LYS:O	1:B:370:ILE:HA	2.20	0.42
1:A:36:ALA:HB2	1:A:459:TRP:CE2	2.54	0.42
1:A:322:GLN:H	1:A:322:GLN:NE2	2.17	0.42
1:C:341:ILE:HD13	1:C:357:PHE:CE2	2.55	0.42
1:A:278:LEU:O	1:A:282:ARG:N	2.52	0.42
1:A:187:TRP:CG	1:C:456:GLN:HE21	2.39	0.41
1:A:191:LEU:O	1:A:192:ILE:HB	2.20	0.41
1:A:361:ARG:O	1:A:433:ILE:HG23	2.20	0.41
1:A:291:ASP:OD1	1:A:291:ASP:N	2.53	0.41
1:A:84:HIS:CG	1:A:85:PRO:HD2	2.55	0.41
1:A:418:ASN:ND2	1:A:444:ALA:HB1	2.36	0.41
1:B:202:VAL:HG11	1:B:239:LYS:HG3	2.02	0.41
1:C:37:LEU:O	1:C:41:SER:OG	2.38	0.41
1:B:408:LEU:O	1:B:412:MET:HG2	2.21	0.41
1:C:55:ASN:OD1	1:C:55:ASN:C	2.63	0.41
1:C:110:MET:HE3	1:C:110:MET:HB2	2.02	0.40
1:A:138:THR:HA	1:A:172:LEU:HD11	2.04	0.40
1:C:94:PHE:CD1	1:C:94:PHE:C	2.99	0.40
1:B:45:ARG:NH1	1:B:69:GLN:OE1	2.46	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	457/460 (99%)	430 (94%)	24 (5%)	3 (1%)	18	48
1	B	457/460 (99%)	431 (94%)	23 (5%)	3 (1%)	18	48
1	C	459/460 (100%)	436 (95%)	21 (5%)	2 (0%)	30	59
All	All	1373/1380 (100%)	1297 (94%)	68 (5%)	8 (1%)	21	51

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	192	ILE
1	B	192	ILE
1	C	192	ILE
1	A	220	ALA
1	C	80	GLY
1	A	286	ALA
1	B	293	ALA
1	B	311	GLY

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	336/351 (96%)	322 (96%)	14 (4%)	26	60
1	B	349/351 (99%)	337 (97%)	12 (3%)	32	66
1	C	349/351 (99%)	333 (95%)	16 (5%)	24	57

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1034/1053 (98%)	992 (96%)	42 (4%)	27 61

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	19	SER
1	A	28	GLN
1	A	60	VAL
1	A	64	ARG
1	A	94	PHE
1	A	160	ARG
1	A	214	ASN
1	A	229	SER
1	A	288	LEU
1	A	291	ASP
1	A	308	VAL
1	A	322	GLN
1	A	341	ILE
1	A	394	SER
1	B	48	GLN
1	B	64	ARG
1	B	94	PHE
1	B	171	LYS
1	B	193	ARG
1	B	258	ILE
1	B	262	THR
1	B	304	GLN
1	B	315	VAL
1	B	386	SER
1	B	414	SER
1	B	431	ASP
1	C	1	MET
1	C	17	LEU
1	C	41	SER
1	C	68	VAL
1	C	70	PHE
1	C	94	PHE
1	C	127	SER
1	C	175	SER
1	C	186	SER
1	C	206	GLU
1	C	229	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	238	LEU
1	C	260	LYS
1	C	313	VAL
1	C	341	ILE
1	C	386	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	61	GLN
1	A	254	GLN
1	A	322	GLN
1	B	48	GLN
1	B	207	HIS
1	C	99	GLN
1	C	207	HIS
1	C	284	GLN
1	C	351	GLN
1	C	392	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The

Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NDP	B	501	-	51,52,52	1.28	5 (9%)	71,80,80	1.84	15 (21%)
4	TLA	A	507	-	9,9,9	1.14	0	12,12,12	1.50	2 (16%)
3	GOL	A	506	-	5,5,5	0.32	0	5,5,5	0.19	0
3	GOL	B	502	-	5,5,5	0.28	0	5,5,5	0.31	0
3	GOL	C	504	-	5,5,5	0.66	0	5,5,5	0.52	0
3	GOL	C	503	-	5,5,5	0.31	0	5,5,5	0.31	0
3	GOL	C	507	-	5,5,5	0.51	0	5,5,5	0.30	0
3	GOL	A	502	-	5,5,5	0.58	0	5,5,5	0.69	0
3	GOL	C	505	-	5,5,5	0.73	0	5,5,5	0.96	0
3	GOL	C	502	-	5,5,5	0.46	0	5,5,5	0.32	0
3	GOL	A	505	-	5,5,5	0.75	0	5,5,5	0.63	0
3	GOL	C	506	-	5,5,5	0.69	0	5,5,5	0.53	0
4	TLA	B	503	-	9,9,9	1.30	0	12,12,12	1.48	3 (25%)
2	NDP	C	501	-	51,52,52	1.22	3 (5%)	71,80,80	1.67	16 (22%)
4	TLA	C	508	-	9,9,9	1.47	2 (22%)	12,12,12	1.92	3 (25%)
2	NDP	A	501	-	51,52,52	1.49	7 (13%)	71,80,80	1.65	14 (19%)
3	GOL	A	504	-	5,5,5	0.18	0	5,5,5	0.55	0
3	GOL	A	503	-	5,5,5	0.50	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NDP	B	501	-	-	8/34/77/77	0/5/5/5
4	TLA	A	507	-	-	5/12/12/12	-
3	GOL	A	506	-	-	1/4/4/4	-
3	GOL	B	502	-	-	2/4/4/4	-
3	GOL	C	504	-	-	0/4/4/4	-
3	GOL	C	503	-	-	0/4/4/4	-
3	GOL	C	507	-	-	1/4/4/4	-
3	GOL	A	502	-	-	2/4/4/4	-
3	GOL	C	505	-	-	3/4/4/4	-
3	GOL	C	502	-	-	2/4/4/4	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	505	-	-	1/4/4/4	-
3	GOL	C	506	-	-	4/4/4/4	-
4	TLA	B	503	-	-	12/12/12/12	-
2	NDP	C	501	-	-	5/34/77/77	0/5/5/5
4	TLA	C	508	-	-	10/12/12/12	-
2	NDP	A	501	-	-	5/34/77/77	0/5/5/5
3	GOL	A	504	-	-	0/4/4/4	-
3	GOL	A	503	-	-	2/4/4/4	-

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	501	NDP	C5A-C4A	5.70	1.49	1.39
2	B	501	NDP	C5A-C4A	4.90	1.47	1.39
2	C	501	NDP	C5A-C4A	4.83	1.47	1.39
2	A	501	NDP	PA-O3	3.63	1.63	1.59
2	B	501	NDP	C5A-N7A	-3.54	1.32	1.39
2	A	501	NDP	C8A-N7A	2.74	1.36	1.31
2	B	501	NDP	C4A-N9A	-2.61	1.32	1.37
4	C	508	TLA	C2-C1	-2.60	1.48	1.52
2	C	501	NDP	C5A-N7A	-2.52	1.34	1.39
2	A	501	NDP	C5A-C6A	2.41	1.47	1.41
2	A	501	NDP	C4A-N9A	-2.34	1.32	1.37
2	C	501	NDP	C8A-N7A	2.29	1.36	1.31
2	B	501	NDP	P2B-O2B	2.26	1.63	1.59
2	A	501	NDP	C2N-C3N	2.21	1.41	1.35
2	A	501	NDP	PN-O3	2.19	1.61	1.59
4	C	508	TLA	O41-C4	-2.17	1.23	1.30
2	B	501	NDP	C8A-N7A	2.09	1.35	1.31

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	501	NDP	O4D-C1D-N1N	5.44	118.46	108.08
2	B	501	NDP	C5A-C4A-N3A	-5.23	119.51	126.72
2	C	501	NDP	N3A-C4A-N9A	5.16	135.95	127.17
2	B	501	NDP	C2B-C1B-N9A	-5.09	105.38	113.75
2	A	501	NDP	C4A-N9A-C8A	4.95	110.94	105.74
2	B	501	NDP	N3A-C4A-N9A	4.84	135.40	127.17
2	C	501	NDP	C5A-C4A-N3A	-4.77	120.14	126.72
4	C	508	TLA	O2-C2-C1	-4.68	100.69	110.69

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	501	NDP	N3A-C4A-N9A	4.65	135.07	127.17
2	C	501	NDP	C4A-N9A-C8A	4.18	110.13	105.74
2	A	501	NDP	C5A-C4A-N3A	-4.05	121.14	126.72
2	B	501	NDP	N3A-C2A-N1A	-3.94	122.62	128.58
2	C	501	NDP	O2B-P2B-O1X	-3.62	96.42	109.33
2	B	501	NDP	C2A-N3A-C4A	3.43	120.22	111.83
2	C	501	NDP	N3A-C2A-N1A	-3.43	123.39	128.58
2	B	501	NDP	C3N-C7N-N7N	3.12	123.21	117.67
2	C	501	NDP	C6N-N1N-C2N	2.99	122.52	119.32
2	C	501	NDP	C2A-N3A-C4A	2.98	119.11	111.83
4	B	503	TLA	O41-C4-C3	2.98	121.58	113.31
4	C	508	TLA	C3-C2-C1	2.96	116.39	109.82
2	C	501	NDP	N6A-C6A-N1A	2.94	124.92	118.38
2	A	501	NDP	C2B-C1B-N9A	-2.88	109.01	113.75
2	A	501	NDP	O3X-P2B-O2X	2.84	118.45	107.80
4	A	507	TLA	C3-C2-C1	2.76	115.94	109.82
2	B	501	NDP	N6A-C6A-N1A	2.75	124.51	118.38
2	A	501	NDP	P2B-O2B-C2B	-2.72	116.16	123.43
2	A	501	NDP	N6A-C6A-N1A	2.71	124.41	118.38
2	B	501	NDP	O4B-C1B-N9A	2.67	113.22	108.09
2	C	501	NDP	C5A-C6A-N6A	-2.65	116.72	123.29
2	B	501	NDP	O3-PA-O1A	-2.60	102.89	110.70
2	B	501	NDP	P2B-O2B-C2B	-2.55	116.62	123.43
2	A	501	NDP	C2A-N3A-C4A	2.50	117.94	111.83
2	A	501	NDP	C1D-N1N-C6N	-2.46	115.57	120.77
2	A	501	NDP	N3A-C2A-N1A	-2.44	124.89	128.58
4	B	503	TLA	O41-C4-O4	-2.34	118.76	124.08
2	B	501	NDP	C5A-C6A-N6A	-2.34	117.50	123.29
4	A	507	TLA	O41-C4-C3	2.33	119.79	113.31
2	C	501	NDP	C1D-N1N-C6N	-2.33	115.85	120.77
2	A	501	NDP	C5A-C6A-N6A	-2.32	117.54	123.29
2	A	501	NDP	O5B-PA-O1A	-2.29	99.85	108.94
2	C	501	NDP	N9A-C8A-N7A	-2.28	110.70	113.94
2	C	501	NDP	O2N-PN-O1N	2.27	123.01	112.44
2	C	501	NDP	P2B-O2B-C2B	-2.27	117.37	123.43
2	A	501	NDP	N9A-C8A-N7A	-2.22	110.79	113.94
2	A	501	NDP	C6N-N1N-C2N	2.22	121.69	119.32
2	C	501	NDP	O4B-C4B-C3B	2.17	109.45	105.15
2	B	501	NDP	O2A-PA-O1A	2.16	122.47	112.44
2	C	501	NDP	O3X-P2B-O2X	2.14	115.83	107.80
2	B	501	NDP	C2A-N1A-C6A	2.11	122.19	118.73
2	C	501	NDP	O4B-C4B-C5B	-2.06	102.72	109.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	503	TLA	C3-C2-C1	2.06	114.38	109.82
4	C	508	TLA	O1-C1-C2	-2.04	116.18	121.62
2	B	501	NDP	O2D-C2D-C3D	2.02	118.30	111.82

There are no chirality outliers.

All (63) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	B	501	NDP	C5B-O5B-PA-O3
3	A	502	GOL	O1-C1-C2-C3
3	A	503	GOL	C1-C2-C3-O3
3	B	502	GOL	O1-C1-C2-C3
3	C	505	GOL	O1-C1-C2-C3
3	C	506	GOL	C1-C2-C3-O3
3	C	506	GOL	O2-C2-C3-O3
4	B	503	TLA	C1-C2-C3-O3
4	B	503	TLA	O2-C2-C3-O3
4	B	503	TLA	O2-C2-C3-C4
4	B	503	TLA	C1-C2-C3-C4
4	C	508	TLA	O1-C1-C2-O2
4	C	508	TLA	O11-C1-C2-O2
4	B	503	TLA	O11-C1-C2-C3
4	B	503	TLA	C2-C3-C4-O4
4	B	503	TLA	C2-C3-C4-O41
2	C	501	NDP	O4D-C4D-C5D-O5D
4	B	503	TLA	O3-C3-C4-O4
4	A	507	TLA	O2-C2-C3-O3
4	A	507	TLA	O2-C2-C3-C4
4	B	503	TLA	O3-C3-C4-O41
4	A	507	TLA	C1-C2-C3-O3
4	C	508	TLA	O3-C3-C4-O4
4	B	503	TLA	O1-C1-C2-C3
2	B	501	NDP	C3B-C2B-O2B-P2B
3	C	502	GOL	C1-C2-C3-O3
3	C	506	GOL	O1-C1-C2-C3
4	C	508	TLA	O3-C3-C4-O41
4	C	508	TLA	O11-C1-C2-C3
3	A	502	GOL	O1-C1-C2-O2
3	B	502	GOL	O1-C1-C2-O2
3	C	505	GOL	O1-C1-C2-O2
2	C	501	NDP	O4D-C1D-N1N-C6N
4	C	508	TLA	O1-C1-C2-C3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
3	A	503	GOL	O2-C2-C3-O3
2	C	501	NDP	C3D-C4D-C5D-O5D
4	C	508	TLA	C1-C2-C3-O3
4	A	507	TLA	C1-C2-C3-C4
2	B	501	NDP	PN-O3-PA-O1A
3	A	506	GOL	O2-C2-C3-O3
4	B	503	TLA	O1-C1-C2-O2
2	B	501	NDP	O4D-C1D-N1N-C6N
3	C	505	GOL	C1-C2-C3-O3
4	B	503	TLA	O11-C1-C2-O2
4	C	508	TLA	C2-C3-C4-O4
4	C	508	TLA	C2-C3-C4-O41
2	A	501	NDP	O4D-C1D-N1N-C6N
2	B	501	NDP	C5B-O5B-PA-O1A
3	C	506	GOL	O1-C1-C2-O2
2	B	501	NDP	PN-O3-PA-O2A
2	B	501	NDP	C1B-C2B-O2B-P2B
2	A	501	NDP	O4B-C4B-C5B-O5B
3	C	507	GOL	C1-C2-C3-O3
2	A	501	NDP	C3B-C4B-C5B-O5B
2	A	501	NDP	C2N-C3N-C7N-N7N
2	B	501	NDP	C2N-C3N-C7N-N7N
2	C	501	NDP	C2N-C3N-C7N-N7N
4	C	508	TLA	C1-C2-C3-C4
3	A	505	GOL	O2-C2-C3-O3
3	C	502	GOL	O2-C2-C3-O3
2	C	501	NDP	O4B-C4B-C5B-O5B
2	A	501	NDP	C2B-O2B-P2B-O2X
4	A	507	TLA	O11-C1-C2-C3


There are no ring outliers.


2 monomers are involved in 1 short contact:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	501	NDP	1	0
3	A	504	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier.


Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	459/460 (99%)	0.18	2 (0%) 88 85	27, 42, 56, 75	1 (0%)
1	B	459/460 (99%)	0.19	2 (0%) 88 85	29, 42, 59, 74	1 (0%)
1	C	460/460 (100%)	0.01	2 (0%) 88 85	17, 40, 53, 76	1 (0%)
All	All	1378/1380 (99%)	0.13	6 (0%) 88 85	17, 41, 56, 76	3 (0%)

All (6) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1	MET	4.4
1	B	293	ALA	3.5
1	A	293	ALA	2.7
1	B	312	ALA	2.5
1	A	264	LYS	2.1
1	C	289	VAL	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	GOL	C	507	6/6	0.78	0.16	58,60,62,63	0
3	GOL	A	505	6/6	0.79	0.23	53,54,56,57	0
3	GOL	C	505	6/6	0.85	0.18	46,51,51,51	0
3	GOL	C	504	6/6	0.85	0.20	52,60,63,65	0
3	GOL	A	502	6/6	0.86	0.15	52,56,56,56	0
3	GOL	A	504	6/6	0.87	0.15	56,59,62,66	0
4	TLA	B	503	10/10	0.87	0.15	51,55,60,63	0
3	GOL	A	506	6/6	0.89	0.15	53,56,58,59	0
3	GOL	C	506	6/6	0.89	0.16	46,47,49,51	0
4	TLA	C	508	10/10	0.89	0.11	44,52,55,62	0
4	TLA	A	507	10/10	0.90	0.11	31,37,40,46	0
3	GOL	C	502	6/6	0.92	0.11	41,43,44,44	0
3	GOL	A	503	6/6	0.92	0.13	40,46,52,54	0
3	GOL	B	502	6/6	0.94	0.10	37,40,41,42	0
3	GOL	C	503	6/6	0.95	0.10	48,49,50,53	0
2	NDP	B	501	48/48	0.95	0.09	28,37,46,50	0
2	NDP	A	501	48/48	0.96	0.08	27,32,44,54	0
2	NDP	C	501	48/48	0.97	0.08	30,36,51,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around TLA B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around TLA C 508:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around TLA A 507:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)







Electron density around NDP B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NDP A 501:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.