



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 11:17 AM UTC

PDB ID : 2EFD / pdb_00002efd
Title : Ara7/AtVps9a
Authors : Uejima, T.; Ihara, K.; Wakatsuki, S.
Deposited on : 2007-02-22
Resolution : 3.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION





The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2672 (3.00-3.00)
Clashscore	190562	2977 (3.00-3.00)
Ramachandran outliers	187476	2877 (3.00-3.00)
Sidechain outliers	187428	2880 (3.00-3.00)
RSRZ outliers	180081	2671 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	 63% 25% • 8%
1	C	267	 73% 16% • 8%
2	B	181	 59% 18% 7% 16%
2	D	181	 57% 25% • 16%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6266 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Similarity to vacuolar protein sorting-associated protein VPS9.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	246	Total	C	N	O	S	0	0	0
			1953	1242	324	375	12			
1	C	246	Total	C	N	O	S	0	0	0
			1953	1242	324	375	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP Q9LT31
A	0	SER	-	expression tag	UNP Q9LT31
C	-1	GLY	-	expression tag	UNP Q9LT31
C	0	SER	-	expression tag	UNP Q9LT31

- Molecule 2 is a protein called Small GTP-binding protein-like.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	152	Total	C	N	O	S	0	0	0
			1180	752	202	222	4			
2	D	152	Total	C	N	O	S	0	0	0
			1180	752	202	222	4			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-1	GLY	-	expression tag	UNP Q9LT31
B	0	SER	-	expression tag	UNP Q9LT31
D	-1	GLY	-	expression tag	UNP Q9LT31
D	0	SER	-	expression tag	UNP Q9LT31


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Similarity to vacuolar protein sorting-associated protein VPS9



- Molecule 1: Similarity to vacuolar protein sorting-associated protein VPS9




- Molecule 2: Small GTP-binding protein-like



- Molecule 2: Small GTP-binding protein-like





4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	189.54Å 189.54Å 75.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	38.95 – 3.00 38.95 – 3.00	Depositor EDS
% Data completeness (in resolution range)	95.9 (38.95-3.00) 95.9 (38.95-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.23 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.306 , 0.351 0.304 , 0.347	Depositor DCC
R_{free} test set	1494 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	74.0	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 84.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	0.076 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.89	EDS
Total number of atoms	6266	wwPDB-VP
Average B, all atoms (Å ²)	74.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.17% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.57	0/1991	0.94	2/2687 (0.1%)
1	C	0.54	0/1991	0.96	3/2687 (0.1%)
2	B	0.55	0/1200	0.92	3/1622 (0.2%)
2	D	0.54	0/1200	0.90	0/1622
All	All	0.55	0/6382	0.93	8/8618 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	68	SER	N-CA-C	6.65	118.41	111.03
1	C	42	ASP	CA-C-N	5.95	127.28	119.84
1	C	42	ASP	C-N-CA	5.95	127.28	119.84
1	A	178	ASN	N-CA-C	5.89	120.27	113.38
2	B	113	ASN	CA-C-N	5.72	126.99	119.84
2	B	113	ASN	C-N-CA	5.72	126.99	119.84
2	B	95	GLN	N-CA-C	5.33	122.16	110.80
1	C	246	SER	N-CA-C	5.11	116.89	110.91

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1953	0	1937	41	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1953	0	1937	30	0
2	B	1180	0	1174	20	0
2	D	1180	0	1174	22	0
All	All	6266	0	6222	107	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (107) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:105:VAL:HG12	2:D:109:GLN:HE21	1.42	0.83
1:C:128:PRO:HA	1:C:131:GLN:HG3	1.63	0.79
1:A:48:ALA:O	1:A:52:GLU:HB2	1.87	0.73
1:C:108:LEU:HG	1:C:112:MET:HE2	1.71	0.73
1:C:111:LYS:O	1:C:115:VAL:HG23	1.89	0.71
1:A:41:PRO:HA	1:A:93:ARG:HH21	1.55	0.71
2:D:105:VAL:HG12	2:D:109:GLN:NE2	2.08	0.68
1:A:84:LYS:HG3	1:A:151:ALA:HB2	1.75	0.68
1:A:176:ALA:HB2	1:C:138:LEU:HD21	1.79	0.65
1:A:161:ASN:O	1:A:165:VAL:HG23	1.99	0.63
1:A:64:HIS:CE1	1:A:66:LEU:HB2	2.34	0.63
2:B:51:THR:HA	2:B:60:LYS:HA	1.80	0.62
1:A:38:ASN:O	1:A:38:ASN:CG	2.41	0.62
1:C:115:VAL:HA	1:C:118:PHE:CE2	2.35	0.61
2:B:29:ARG:HH11	2:B:157:ALA:HB2	1.67	0.59
1:C:128:PRO:CA	1:C:131:GLN:HG3	2.33	0.59
1:C:170:LEU:HD21	1:C:190:VAL:HG21	1.83	0.59
2:D:120:LEU:HD23	2:D:149:PHE:HD2	1.66	0.59
1:A:156:LEU:HD23	1:A:160:LEU:HG	1.84	0.58
2:D:163:ILE:O	2:D:167:ILE:HG13	2.03	0.58
2:B:61:PHE:HB3	2:B:63:ILE:HD11	1.86	0.57
2:D:107:GLU:O	2:D:111:GLN:HG2	2.04	0.57
1:C:229:ASN:O	1:C:232:SER:OG	2.17	0.57
1:C:83:GLU:CD	1:C:152:PRO:HG2	2.31	0.56
1:C:146:ILE:HG23	1:C:147:ASN:OD1	2.04	0.56
1:A:241:ASP:O	1:A:242:ALA:C	2.49	0.56
2:B:166:GLU:O	2:B:170:ARG:HG3	2.07	0.54
1:A:105:ASP:OD2	1:A:204:HIS:HB3	2.08	0.54
1:A:39:ASN:HB2	1:A:93:ARG:HH12	1.73	0.54
1:A:41:PRO:HA	1:A:93:ARG:NH2	2.22	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:167:ASN:O	1:A:171:LEU:HB2	2.09	0.53
2:B:83:ALA:HB3	2:B:116:MET:HE2	1.89	0.53
1:A:157:VAL:O	1:A:158:CYS:C	2.52	0.53
1:C:226:PHE:O	1:C:229:ASN:HB2	2.09	0.52
2:D:17:GLY:O	2:D:23:LYS:HE3	2.10	0.52
2:B:126:ASP:OD2	2:B:126:ASP:N	2.39	0.52
1:A:108:LEU:HD21	1:A:234:GLU:HG3	1.92	0.52
1:A:115:VAL:HA	1:A:118:PHE:CE2	2.45	0.51
1:C:84:LYS:HG3	1:C:151:ALA:HB2	1.91	0.51
1:A:225:TYR:OH	2:B:65:ASP:O	2.29	0.50
1:A:36:PHE:O	1:A:93:ARG:NH1	2.45	0.50
1:C:153:ARG:O	1:C:157:VAL:HG23	2.11	0.50
1:C:242:ALA:HB1	1:C:247:LEU:O	2.10	0.50
2:B:13:LEU:HD21	2:B:87:ILE:HG13	1.93	0.50
2:D:91:ASP:HB3	2:D:94:ASN:HB2	1.93	0.50
1:A:192:ILE:O	1:A:193:TYR:C	2.56	0.49
2:B:94:ASN:O	2:B:96:ALA:N	2.46	0.49
1:C:128:PRO:HA	1:C:131:GLN:CG	2.41	0.48
2:D:120:LEU:HD23	2:D:149:PHE:CD2	2.47	0.48
1:C:132:ASN:HB3	1:C:136:TRP:NE1	2.28	0.48
1:A:163:CYS:SG	1:A:187:PHE:HZ	2.37	0.47
1:A:37:SER:HA	1:A:93:ARG:HD2	1.96	0.47
1:A:44:GLU:H	1:A:44:GLU:CD	2.21	0.47
1:A:199:ASN:O	1:A:201:PRO:HD3	2.13	0.47
1:C:214:ARG:O	1:C:215:ARG:C	2.58	0.47
1:C:259:ALA:C	1:C:261:ALA:H	2.21	0.47
1:A:122:GLU:O	1:A:123:ASN:C	2.56	0.47
1:A:138:LEU:HB3	1:A:165:VAL:HG11	1.97	0.46
2:D:87:ILE:HA	2:D:119:ALA:O	2.16	0.46
2:B:61:PHE:HB3	2:B:63:ILE:CD1	2.45	0.46
2:D:90:PHE:HB2	2:D:97:SER:HB2	1.97	0.46
1:C:120:SER:OG	1:C:122:GLU:HG2	2.16	0.46
2:D:76:ALA:N	2:D:77:PRO:HD2	2.31	0.46
2:D:77:PRO:HA	2:D:80:TYR:CE1	2.51	0.46
1:C:185:ASP:O	1:C:189:PRO:HG2	2.16	0.45
1:C:125:ASP:CG	2:D:71:ARG:HE	2.24	0.45
1:C:126:ILE:HG22	1:C:131:GLN:HG2	1.98	0.45
1:A:87:MET:HE2	1:A:213:TYR:CG	2.52	0.45
1:C:124:LEU:HB3	2:D:72:TYR:OH	2.17	0.45
1:A:61:PHE:HE1	1:A:82:LEU:HD22	1.81	0.44
1:C:132:ASN:HB3	1:C:136:TRP:CE2	2.52	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:24:SER:O	2:B:28:LEU:HD23	2.18	0.44
1:C:107:LYS:NZ	1:C:111:LYS:HE3	2.32	0.44
1:A:140:GLN:OE1	1:A:197:LYS:HB3	2.17	0.44
2:B:8:SER:HB3	2:B:58:THR:O	2.17	0.44
1:A:90:LEU:O	1:A:94:VAL:HG22	2.18	0.43
1:A:29:ILE:O	1:A:33:ILE:HG13	2.19	0.43
1:A:55:SER:O	1:A:56:LYS:C	2.60	0.43
1:C:132:ASN:OD1	1:C:132:ASN:C	2.62	0.43
2:B:9:ILE:HB	2:B:59:VAL:HG23	2.01	0.43
2:D:23:LYS:HB3	2:D:89:VAL:HG21	2.01	0.43
1:C:192:ILE:O	1:C:196:ILE:HG13	2.19	0.43
2:D:140:THR:O	2:D:144:GLU:HG3	2.19	0.42
2:B:54:VAL:HG12	2:B:165:TYR:CD2	2.54	0.42
2:B:168:ALA:HA	2:B:171:LEU:HD12	2.01	0.42
1:A:64:HIS:HE1	1:A:66:LEU:HB2	1.82	0.42
1:A:39:ASN:O	1:A:93:ARG:CZ	2.67	0.42
2:B:152:THR:HA	2:B:159:ASN:HB2	2.02	0.42
2:D:90:PHE:CE1	2:D:98:PHE:HB2	2.54	0.42
1:A:73:GLU:H	1:A:73:GLU:HG3	1.60	0.42
1:C:71:SER:O	1:C:75:LEU:HG	2.18	0.42
2:D:160:VAL:O	2:D:163:ILE:HG22	2.20	0.42
1:A:222:GLU:O	1:A:226:PHE:HD1	2.03	0.42
1:A:160:LEU:O	1:A:164:LYS:HG3	2.19	0.41
2:B:30:PHE:CE1	2:B:50:GLN:HG2	2.56	0.41
1:C:189:PRO:HG3	2:D:72:TYR:CE1	2.55	0.41
2:D:113:ASN:HA	2:D:114:PRO:HD3	1.82	0.41
1:A:19:LYS:C	1:A:21:SER:H	2.28	0.41
1:C:120:SER:O	1:C:123:ASN:HB2	2.21	0.41
1:A:19:LYS:HA	1:A:20:PRO:HD3	1.95	0.41
2:D:94:ASN:O	2:D:95:GLN:C	2.64	0.40
1:A:39:ASN:O	1:A:93:ARG:NH1	2.54	0.40
2:D:165:TYR:O	2:D:168:ALA:HB3	2.21	0.40
2:B:113:ASN:HA	2:B:114:PRO:HD3	1.84	0.40
1:A:63:ALA:O	1:A:64:HIS:C	2.64	0.40
1:A:232:SER:HB2	2:B:75:LEU:HD13	2.04	0.40
2:B:171:LEU:HA	2:B:172:PRO:HD2	1.83	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	244/267 (91%)	209 (86%)	31 (13%)	4 (2%)	7	34
1	C	244/267 (91%)	218 (89%)	23 (9%)	3 (1%)	10	40
2	B	148/181 (82%)	132 (89%)	13 (9%)	3 (2%)	6	28
2	D	148/181 (82%)	123 (83%)	20 (14%)	5 (3%)	3	17
All	All	784/896 (88%)	682 (87%)	87 (11%)	15 (2%)	6	30

All (15) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	67	TRP
2	B	95	GLN
2	D	56	ASP
1	A	215	ARG
2	D	124	LYS
2	B	56	ASP
1	C	215	ARG
2	D	95	GLN
2	B	157	ALA
2	D	74	SER
2	D	154	ALA
1	A	20	PRO
1	A	71	SER
1	C	37	SER
1	C	43	PRO

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	217/235 (92%)	195 (90%)	22 (10%)	7	29
1	C	217/235 (92%)	206 (95%)	11 (5%)	21	55
2	B	120/143 (84%)	101 (84%)	19 (16%)	2	13
2	D	120/143 (84%)	106 (88%)	14 (12%)	5	23
All	All	674/756 (89%)	608 (90%)	66 (10%)	7	30

All (66) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	30	LYS
1	A	34	VAL
1	A	38	ASN
1	A	44	GLU
1	A	56	LYS
1	A	62	ARG
1	A	67	TRP
1	A	73	GLU
1	A	93	ARG
1	A	99	THR
1	A	108	LEU
1	A	122	GLU
1	A	137	LEU
1	A	171	LEU
1	A	179	GLU
1	A	194	VAL
1	A	197	LYS
1	A	214	ARG
1	A	228	THR
1	A	231	LEU
1	A	238	SER
1	A	254	LYS
2	B	9	ILE
2	B	13	LEU
2	B	18	ASP
2	B	24	SER
2	B	27	VAL
2	B	28	LEU
2	B	49	SER
2	B	50	GLN
2	B	51	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	54	VAL
2	B	56	ASP
2	B	60	LYS
2	B	62	GLU
2	B	100	ARG
2	B	115	ASN
2	B	126	ASP
2	B	128	LEU
2	B	155	LYS
2	B	158	THR
1	C	37	SER
1	C	56	LYS
1	C	111	LYS
1	C	122	GLU
1	C	132	ASN
1	C	156	LEU
1	C	197	LYS
1	C	214	ARG
1	C	231	LEU
1	C	246	SER
1	C	254	LYS
2	D	9	ILE
2	D	10	ASN
2	D	13	LEU
2	D	28	LEU
2	D	54	VAL
2	D	58	THR
2	D	60	LYS
2	D	62	GLU
2	D	89	VAL
2	D	100	ARG
2	D	127	LEU
2	D	150	MET
2	D	153	SER
2	D	158	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	116	GLN
1	A	167	ASN
1	A	204	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	94	ASN
2	B	109	GLN
1	C	116	GLN
1	C	117	GLN
1	C	202	GLN
2	D	10	ASN
2	D	94	ASN
2	D	109	GLN
2	D	139	GLN
2	D	143	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å²)	Q<0.9	
1	A	246/267 (92%)	-1.15	0	100	100	43, 56, 94, 101	0
1	C	246/267 (92%)	-1.05	0	100	100	49, 65, 114, 127	0
2	B	152/181 (83%)	-0.89	0	100	100	46, 86, 106, 109	0
2	D	152/181 (83%)	-0.78	1 (0%)	84	66	65, 92, 116, 118	0
All	All	796/896 (88%)	-1.00	1 (0%)	92	86	43, 69, 113, 127	0

All (1) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	53	ALA	4.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.