



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:14 AM UTC

PDB ID : 8EMB / pdb\_00008emb  
Title : X-ray crystal structure of Thermosynechococcus elongatus Si3 domain of RNA polymerase RpoC2 subunit  
Authors : Murakami, K.S.; Imashimizu, M.; Qayyum, M.Z.; Vishwakarma, R.K.; Yuzenkova, Y.  
Deposited on : 2022-09-27  
Resolution : 3.06 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.06 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	2469 (3.10-3.02)
Clashscore	190562	2569 (3.10-3.02)
Ramachandran outliers	187476	2424 (3.10-3.02)
Sidechain outliers	187428	2423 (3.10-3.02)
RSRZ outliers	180081	2469 (3.10-3.02)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	<div> <div>3%</div> <div>76%</div> <div>23%</div> <div>.</div> </div>
1	B	553	<div> <div>4%</div> <div>77%</div> <div>20%</div> <div>.</div> </div>
1	C	553	<div> <div>2%</div> <div>71%</div> <div>25%</div> <div>.</div> </div>
1	D	553	<div> <div>5%</div> <div>72%</div> <div>26%</div> <div>.</div> </div>
1	E	553	<div> <div>9%</div> <div>44%</div> <div>21%</div> <div>35%</div> </div>

*Continued on next page...*

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	553	<div><div><div></div><div></div><div></div><div></div></div><div>2%34%10%56%</div></div>

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 21086 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	550	Total	C	N	O	S	Se	0	0	0
			4185	2609	736	832	2	6			
1	B	541	Total	C	N	O	S	Se	0	0	0
			4127	2574	726	819	2	6			
1	C	531	Total	C	N	O	S	Se	0	0	0
			4047	2526	711	803	2	5			
1	D	543	Total	C	N	O	S	Se	0	0	0
			4140	2582	728	823	2	5			
1	E	358	Total	C	N	O	S	Se	0	0	0
			2752	1718	476	554	2	2			
1	F	246	Total	C	N	O	Se		0	0	0
			1835	1134	335	363	3				

There are 42 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	431	GLY	-	expression tag	UNP Q8DL57
A	432	SER	-	expression tag	UNP Q8DL57
A	433	HIS	-	expression tag	UNP Q8DL57
A	434	MSE	-	expression tag	UNP Q8DL57
A	508	MSE	LEU	engineered mutation	UNP Q8DL57
A	738	MSE	LEU	engineered mutation	UNP Q8DL57
A	922	MSE	LEU	engineered mutation	UNP Q8DL57
B	431	GLY	-	expression tag	UNP Q8DL57
B	432	SER	-	expression tag	UNP Q8DL57
B	433	HIS	-	expression tag	UNP Q8DL57
B	434	MSE	-	expression tag	UNP Q8DL57
B	508	MSE	LEU	engineered mutation	UNP Q8DL57
B	738	MSE	LEU	engineered mutation	UNP Q8DL57
B	922	MSE	LEU	engineered mutation	UNP Q8DL57
C	431	GLY	-	expression tag	UNP Q8DL57
C	432	SER	-	expression tag	UNP Q8DL57
C	433	HIS	-	expression tag	UNP Q8DL57

*Continued on next page...*


*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
C	434	MSE	-	expression tag	UNP Q8DL57
C	508	MSE	LEU	engineered mutation	UNP Q8DL57
C	738	MSE	LEU	engineered mutation	UNP Q8DL57
C	922	MSE	LEU	engineered mutation	UNP Q8DL57
D	431	GLY	-	expression tag	UNP Q8DL57
D	432	SER	-	expression tag	UNP Q8DL57
D	433	HIS	-	expression tag	UNP Q8DL57
D	434	MSE	-	expression tag	UNP Q8DL57
D	508	MSE	LEU	engineered mutation	UNP Q8DL57
D	738	MSE	LEU	engineered mutation	UNP Q8DL57
D	922	MSE	LEU	engineered mutation	UNP Q8DL57
E	431	GLY	-	expression tag	UNP Q8DL57
E	432	SER	-	expression tag	UNP Q8DL57
E	433	HIS	-	expression tag	UNP Q8DL57
E	434	MSE	-	expression tag	UNP Q8DL57
E	508	MSE	LEU	engineered mutation	UNP Q8DL57
E	738	MSE	LEU	engineered mutation	UNP Q8DL57
E	922	MSE	LEU	engineered mutation	UNP Q8DL57
F	431	GLY	-	expression tag	UNP Q8DL57
F	432	SER	-	expression tag	UNP Q8DL57
F	433	HIS	-	expression tag	UNP Q8DL57
F	434	MSE	-	expression tag	UNP Q8DL57
F	508	MSE	LEU	engineered mutation	UNP Q8DL57
F	738	MSE	LEU	engineered mutation	UNP Q8DL57
F	922	MSE	LEU	engineered mutation	UNP Q8DL57


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA-directed RNA polymerase subunit beta'




- Molecule 1: DNA-directed RNA polymerase subunit beta'




• Molecule 1: DNA-directed RNA polymerase subunit beta'




• Molecule 1: DNA-directed RNA polymerase subunit beta'



• Molecule 1: DNA-directed RNA polymerase subunit beta'







## 4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	178.66Å 178.66Å 281.04Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.61 – 3.06 49.61 – 3.06	Depositor EDS
% Data completeness (in resolution range)	98.1 (49.61-3.06) 87.0 (49.61-3.06)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.62 (at 3.07Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487, PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.264 , 0.284 0.267 , 0.280	Depositor DCC
$R_{free}$ test set	1998 reflections (2.04%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	73.6	Xtriage
Anisotropy	0.236	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 79.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.028 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.89	EDS
Total number of atoms	21086	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	130.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.29% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.18	0/4233	0.44	0/5731
1	B	0.15	0/4173	0.41	0/5647
1	C	0.19	0/4093	0.46	0/5542
1	D	0.15	0/4188	0.44	0/5671
1	E	0.13	0/2787	0.35	0/3777
1	F	0.11	0/1850	0.33	0/2500
All	All	0.16	0/21324	0.42	0/28868

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4185	0	4263	86	0
1	B	4127	0	4207	70	0
1	C	4047	0	4121	99	0
1	D	4140	0	4214	104	0
1	E	2752	0	2781	80	0
1	F	1835	0	1879	36	0
All	All	21086	0	21465	461	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 461 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:828:LEU:HD11	1:E:801:GLU:HG3	1.46	0.96
1:C:588:GLU:HG3	1:C:793:GLN:HG2	1.58	0.86
1:A:815:GLU:HG3	1:A:818:PRO:HD2	1.59	0.84
1:A:589:LEU:HD22	1:A:591:ASP:HB2	1.63	0.80
1:D:587:ALA:HB3	1:D:794:LEU:HD21	1.64	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	548/553 (99%)	517 (94%)	31 (6%)	0	100	100
1	B	537/553 (97%)	504 (94%)	31 (6%)	2 (0%)	30	57
1	C	527/553 (95%)	503 (95%)	24 (5%)	0	100	100
1	D	541/553 (98%)	505 (93%)	36 (7%)	0	100	100
1	E	356/553 (64%)	325 (91%)	31 (9%)	0	100	100
1	F	242/553 (44%)	229 (95%)	13 (5%)	0	100	100
All	All	2751/3318 (83%)	2583 (94%)	166 (6%)	2 (0%)	48	75

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	704	THR
1	B	883	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	456/452 (101%)	453 (99%)	3 (1%)	76	80
1	B	451/452 (100%)	451 (100%)	0	100	100
1	C	442/452 (98%)	441 (100%)	1 (0%)	87	87
1	D	452/452 (100%)	452 (100%)	0	100	100
1	E	305/452 (68%)	304 (100%)	1 (0%)	86	85
1	F	196/452 (43%)	196 (100%)	0	100	100
All	All	2302/2712 (85%)	2297 (100%)	5 (0%)	87	87

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	464	ARG
1	A	467	THR
1	A	469	THR
1	C	902	GLU
1	E	688	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 15 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	763	GLN
1	D	878	GLN
1	C	878	GLN
1	E	595	HIS
1	D	690	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å <sup>2</sup> )	Q<0.9
1	A	544/553 (98%)	0.37	15 (2%)	55	32	57, 122, 177, 218	0
1	B	535/553 (96%)	0.34	24 (4%)	38	20	43, 109, 165, 199	0
1	C	526/553 (95%)	0.19	12 (2%)	61	38	45, 75, 139, 200	0
1	D	538/553 (97%)	0.50	25 (4%)	37	19	58, 144, 192, 216	0
1	E	356/553 (64%)	0.99	48 (13%)	7	4	119, 190, 242, 272	0
1	F	243/553 (43%)	0.63	10 (4%)	41	22	162, 192, 221, 252	0
All	All	2742/3318 (82%)	0.46	134 (4%)	35	17	43, 126, 208, 272	0

The worst 5 of 134 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	808	ASP	5.9
1	A	808	ASP	4.7
1	F	976	VAL	3.9
1	D	977	PHE	3.8
1	B	983	GLY	3.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.