



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 01:35 PM EDT

PDB ID : 8FER / pdb_00008fer
Title : 16mer self-complementary duplex RNA with two continuous s(2)U:s(2)U pairs
Authors : Fang, Z.; Zhou, L.; Szostak, J.W.
Deposited on : 2022-12-06
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)
RNA backbone	3983	1008 (2.10-0.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	16	
1	BBB	16	

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 753 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a RNA chain called RNA 16mer with two continuous s(2)U.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	AAA	16	Total	C	N	O	P	S	0	0	0
			334	151	56	110	15	2			
1	BBB	16	Total	C	N	O	P	S	0	0	0
			334	151	56	110	15	2			

- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	AAA	46	Total	O	0	0
			46	46		
2	BBB	39	Total	O	0	0
			39	39		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: RNA 16mer with two continuous s(2)U



- Molecule 1: RNA 16mer with two continuous s(2)U



4 Data and refinement statistics

Property	Value	Source
Space group	H 3 2	Depositor
Cell constants a, b, c, α , β , γ	43.81Å 43.81Å 258.17Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	43.06 – 1.54 43.06 – 1.54	Depositor EDS
% Data completeness (in resolution range)	96.1 (43.06-1.54) 96.1 (43.06-1.54)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.19 (at 1.54Å)	Xtriage
Refinement program	REFMAC 5.8.0267	Depositor
R, R_{free}	0.219 , 0.259 0.223 , 0.260	Depositor DCC
R_{free} test set	725 reflections (4.90%)	wwPDB-VP
Wilson B-factor (Å ²)	10.5	Xtriage
Anisotropy	0.261	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	753	wwPDB-VP
Average B, all atoms (Å ²)	15.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 44.15 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.5905e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SUR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	AAA	0.87	0/327	1.35	3/505 (0.6%)
1	BBB	0.90	0/327	1.34	1/505 (0.2%)
All	All	0.88	0/654	1.35	4/1010 (0.4%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	AAA	7	G	C4'-C3'-C2'	-5.96	96.64	102.60
1	AAA	1	A	P-O3'-C3'	-5.70	111.65	120.20
1	BBB	4	G	C4'-C3'-C2'	-5.57	97.03	102.60
1	AAA	11	U	C4'-C3'-C2'	-5.04	97.56	102.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	334	0	172	0	0
1	BBB	334	0	172	0	0
2	AAA	46	0	0	0	0
2	BBB	39	0	0	0	0
All	All	753	0	344	0	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 0.

There are no clashes within the asymmetric unit.

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
1	AAA	15/16 (93%)	0	0
1	BBB	15/16 (93%)	0	0
All	All	30/32 (93%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	SUR	AAA	8	1	17,21,22	2.03	4 (23%)	21,30,33	1.60	5 (23%)
1	SUR	BBB	8	1	17,21,22	1.39	4 (23%)	21,30,33	1.58	3 (14%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	SUR	BBB	9	1	17,21,22	2.09	3 (17%)	21,30,33	2.29	6 (28%)
1	SUR	AAA	9	1	17,21,22	1.58	3 (17%)	21,30,33	1.57	2 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	SUR	AAA	8	1	-	0/7/25/26	0/2/2/2
1	SUR	BBB	8	1	-	0/7/25/26	0/2/2/2
1	SUR	BBB	9	1	-	0/7/25/26	0/2/2/2
1	SUR	AAA	9	1	-	0/7/25/26	0/2/2/2

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	BBB	9	SUR	C2-S2	-6.99	1.56	1.67
1	AAA	8	SUR	C2-S2	5.83	1.77	1.67
1	AAA	9	SUR	C2-S2	4.21	1.74	1.67
1	AAA	9	SUR	C4-N3	-3.15	1.33	1.38
1	AAA	8	SUR	O4'-C4'	-3.04	1.38	1.45
1	BBB	8	SUR	C5-C4	-2.93	1.37	1.43
1	BBB	9	SUR	C5-C4	-2.91	1.37	1.43
1	AAA	8	SUR	C5-C4	-2.77	1.37	1.43
1	BBB	8	SUR	C6-C5	2.57	1.41	1.35
1	BBB	8	SUR	O4'-C4'	2.55	1.50	1.45
1	BBB	8	SUR	C2-S2	2.42	1.71	1.67
1	BBB	9	SUR	C6-C5	2.29	1.40	1.35
1	AAA	9	SUR	O4'-C1'	2.17	1.47	1.42
1	AAA	8	SUR	O3'-C3'	2.10	1.48	1.43

All (16) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	9	SUR	O4-C4-C5	-5.75	115.25	125.16
1	AAA	9	SUR	C2-N3-C4	-5.44	120.63	127.33
1	BBB	8	SUR	C2-N3-C4	-5.16	120.98	127.33
1	BBB	9	SUR	C5-C4-N3	4.68	121.36	114.80
1	BBB	9	SUR	C2-N3-C4	-3.89	122.55	127.33
1	AAA	8	SUR	C2-N3-C4	-3.60	122.91	127.33
1	BBB	9	SUR	C6-C5-C4	-3.24	115.39	119.53

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	BBB	9	SUR	O4-C4-N3	2.84	123.39	119.27
1	BBB	8	SUR	O4-C4-C5	-2.62	120.65	125.16
1	AAA	9	SUR	C5-C6-N1	-2.56	117.67	121.84
1	BBB	9	SUR	O3'-C3'-C4'	2.48	118.20	111.08
1	AAA	8	SUR	O4'-C4'-C3'	-2.41	100.37	105.15
1	AAA	8	SUR	C5-C6-N1	-2.25	118.18	121.84
1	AAA	8	SUR	C2'-C3'-C4'	-2.21	98.35	102.61
1	AAA	8	SUR	S2-C2-N3	-2.19	113.98	119.44
1	BBB	8	SUR	O4-C4-N3	2.10	122.32	119.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	AAA	14/16 (87%)	0.96	2 (14%) 6 8	11, 15, 21, 31	0
1	BBB	14/16 (87%)	0.55	0 100 100	10, 15, 18, 18	0
All	All	28/32 (87%)	0.76	2 (7%) 22 27	10, 15, 18, 31	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	AAA	1	A	5.3
1	AAA	2	G	4.0

6.2 Non-standard residues in protein, DNA, RNA chains

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	SUR	AAA	8	20/21	0.96	0.08	11,15,19,20	0
1	SUR	BBB	9	20/21	0.97	0.07	11,12,14,16	0
1	SUR	AAA	9	20/21	0.98	0.06	9,10,12,14	0
1	SUR	BBB	8	20/21	0.98	0.06	9,10,12,14	0

6.3 Carbohydrates

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.