



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 07:31 PM UTC

PDB ID : 7PHV / pdb\_00007phv  
Title : PfCyRPA bound to monoclonal antibody Cy.007 Fab fragment  
Authors : Ragotte, R.J.; Higgins, M.K.  
Deposited on : 2021-08-18  
Resolution : 3.09 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.09 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	343	<div> <div>7%</div> <div>66%</div> <div>22%</div> <div>•</div> <div>8%</div> </div>
1	D	343	<div> <div>10%</div> <div>68%</div> <div>19%</div> <div>•</div> <div>11%</div> </div>
2	B	233	<div> <div>4%</div> <div>81%</div> <div>13%</div> <div>•</div> <div>5%</div> </div>
2	E	233	<div> <div>4%</div> <div>80%</div> <div>13%</div> <div>•</div> <div>6%</div> </div>
3	C	209	<div> <div>6%</div> <div>91%</div> <div>7%</div> <div>•</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
3	F	209	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	NH4	B	301	-	-	-	X

## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 11521 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Cysteine-rich protective antigen.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	316	Total	C	N	O	S	0	0	0
			2633	1697	420	503	13			
1	D	305	Total	C	N	O	S	0	0	0
			2539	1640	404	483	12			

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	147	ALA	SER	conflict	UNP Q8IFM8
A	324	ALA	THR	conflict	UNP Q8IFM8
A	340	ALA	THR	conflict	UNP Q8IFM8
A	363	GLY	-	expression tag	UNP Q8IFM8
A	364	GLY	-	expression tag	UNP Q8IFM8
A	365	GLY	-	expression tag	UNP Q8IFM8
A	366	GLY	-	expression tag	UNP Q8IFM8
A	367	SER	-	expression tag	UNP Q8IFM8
A	368	GLU	-	expression tag	UNP Q8IFM8
A	369	PRO	-	expression tag	UNP Q8IFM8
A	370	GLU	-	expression tag	UNP Q8IFM8
A	371	ALA	-	expression tag	UNP Q8IFM8
D	147	ALA	SER	conflict	UNP Q8IFM8
D	324	ALA	THR	conflict	UNP Q8IFM8
D	340	ALA	THR	conflict	UNP Q8IFM8
D	363	GLY	-	expression tag	UNP Q8IFM8
D	364	GLY	-	expression tag	UNP Q8IFM8
D	365	GLY	-	expression tag	UNP Q8IFM8
D	366	GLY	-	expression tag	UNP Q8IFM8
D	367	SER	-	expression tag	UNP Q8IFM8
D	368	GLU	-	expression tag	UNP Q8IFM8
D	369	PRO	-	expression tag	UNP Q8IFM8
D	370	GLU	-	expression tag	UNP Q8IFM8
D	371	ALA	-	expression tag	UNP Q8IFM8


- Molecule 2 is a protein called Monoclonal antibody Cy.007 heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	221	Total	C	N	O	S	518	0	0
			1619	1013	272	327	7			
2	E	220	Total	C	N	O	S	646	0	0
			1614	1010	271	326	7			

- Molecule 3 is a protein called Monoclonal antibody Cy.007 light chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	207	Total	C	N	O	S	549	0	0
			1557	965	265	323	4			
3	F	207	Total	C	N	O	S	489	0	0
			1557	965	265	323	4			

- Molecule 4 is AMMONIUM ION (CCD ID: NH4) (formula:  $\text{H}_4\text{N}$ ).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	N	0	0
			1	1		
4	E	1	Total	N	0	0
			1	1		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Cysteine-rich protective antigen




- Molecule 1: Cysteine-rich protective antigen





- Molecule 2: Monoclonal antibody Cy.007 heavy chain

Chain B: 





- Molecule 2: Monoclonal antibody Cy.007 heavy chain

Chain E: 





- Molecule 3: Monoclonal antibody Cy.007 light chain

Chain C: 



- Molecule 3: Monoclonal antibody Cy.007 light chain

Chain F: 



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	74.10Å 87.00Å 342.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.24 – 3.09 45.24 – 3.09	Depositor EDS
% Data completeness (in resolution range)	100.0 (45.24-3.09) 99.9 (45.24-3.09)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.04 (at 3.06Å)	Xtriage
Refinement program	BUSTER 2.10.4 (20-OCT-2021)	Depositor
R, $R_{free}$	0.287 , 0.332 0.290 , 0.328	Depositor DCC
$R_{free}$ test set	2060 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	109.4	Xtriage
Anisotropy	0.558	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.31 , 137.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	11521	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	176.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.28% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NH4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.85	0/2692	1.15	11/3634 (0.3%)
1	D	0.78	0/2597	1.07	4/3509 (0.1%)
2	B	0.60	0/1654	0.89	2/2254 (0.1%)
2	E	0.58	0/1649	0.90	0/2247
3	C	0.57	0/1590	0.88	0/2163
3	F	0.58	0/1590	0.86	0/2163
All	All	0.70	0/11772	0.99	17/15970 (0.1%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	LEU	N-CA-C	7.65	121.60	110.28
1	D	287	ASN	CA-CB-CG	6.68	119.28	112.60
1	A	220	LYS	CA-C-N	6.66	130.32	120.87
1	A	220	LYS	C-N-CA	6.66	130.32	120.87
1	A	287	ASN	CA-CB-CG	6.51	119.11	112.60

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2633	0	2528	40	0
1	D	2539	0	2429	36	0
2	B	1619	0	1570	13	0
2	E	1614	0	1565	17	0
3	C	1557	0	1493	5	0
3	F	1557	0	1493	7	0
4	B	1	0	0	1	0
4	E	1	0	0	1	0
All	All	11521	0	11078	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 115 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:325:ALA:HB1	1:D:346:SER:HB2	1.63	0.81
1:A:134:HIS:CE1	1:A:146:ASN:HD22	2.02	0.77
2:B:52:GLU:OE1	4:B:301:NH4:N	2.19	0.75
3:F:108:ARG:HA	3:F:113:GLY:HA2	1.71	0.73
2:E:92:ASP:HB2	2:E:99:ARG:HH12	1.53	0.72

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	308/343 (90%)	289 (94%)	19 (6%)	0	100	100
1	D	295/343 (86%)	270 (92%)	25 (8%)	0	100	100
2	B	217/233 (93%)	207 (95%)	10 (5%)	0	100	100
2	E	216/233 (93%)	204 (94%)	12 (6%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	C	205/209 (98%)	193 (94%)	12 (6%)	0	100	100
3	F	205/209 (98%)	189 (92%)	16 (8%)	0	100	100
All	All	1446/1570 (92%)	1352 (94%)	94 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	297/316 (94%)	260 (88%)	37 (12%)	4	19
1	D	286/316 (90%)	263 (92%)	23 (8%)	11	37
2	B	179/191 (94%)	168 (94%)	11 (6%)	17	46
2	E	179/191 (94%)	169 (94%)	10 (6%)	19	49
3	C	177/179 (99%)	168 (95%)	9 (5%)	21	52
3	F	177/179 (99%)	168 (95%)	9 (5%)	21	52
All	All	1295/1372 (94%)	1196 (92%)	99 (8%)	12	39

5 of 99 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	40	SER
1	D	210	THR
1	D	99	LEU
1	D	154	TYR
1	D	238	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 16 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	277	ASN
1	D	254	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
2	B	202	GLN
1	D	224	GLN
1	A	352	ASN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry ⓘ

Of 2 ligands modelled in this entry, 2 are modelled with single atom - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers ⓘ

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å <sup>2</sup> )	Q<0.9
1	A	316/343 (92%)	0.56	25 (7%)	18 10	82, 109, 149, 160	0
1	D	305/343 (88%)	0.66	33 (10%)	11 6	83, 136, 180, 216	0
2	B	148/233 (63%)	0.55	10 (6%)	23 12	74, 145, 193, 200	1 (0%)
2	E	132/233 (56%)	0.51	9 (6%)	23 12	88, 181, 215, 226	1 (0%)
3	C	136/209 (65%)	0.61	12 (8%)	15 9	159, 187, 240, 246	0
3	F	145/209 (69%)	1.12	26 (17%)	3 1	187, 211, 239, 245	0
All	All	1182/1570 (75%)	0.65	115 (9%)	13 7	74, 149, 222, 246	2 (0%)

The worst 5 of 115 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	F	116	GLY	6.5
3	F	114	ALA	6.2
3	F	106	GLY	6.2
3	F	105	CYS	5.4
3	F	117	ALA	5.2

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	NH4	E	301	1/1	0.46	0.36	118,118,118,118	0
4	NH4	B	301	1/1	0.63	0.53	72,72,72,72	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.