



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 1, 2026 – 07:49 PM UTC

PDB ID : 6PP0 / pdb\_00006pp0  
Title : Structure of human endothelial nitric oxide synthase heme domain in complex with 7-(3-(Aminomethyl)-4-(cyclobutylmethoxy)phenyl)-4-methylquinolin-2-amine  
Authors : Li, H.; Poulos, T.L.  
Deposited on : 2019-07-05  
Resolution : 1.97 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	440	<div> <div>30%</div> <div>71%</div> <div>19%</div> <div>•</div> <div>9%</div> </div>
1	B	440	<div> <div>12%</div> <div>78%</div> <div>13%</div> <div>8%</div> </div>
1	C	440	<div> <div>17%</div> <div>79%</div> <div>11%</div> <div>•</div> <div>9%</div> </div>
1	D	440	<div> <div>15%</div> <div>77%</div> <div>13%</div> <div>•</div> <div>8%</div> </div>

## 2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 13897 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Nitric oxide synthase, endothelial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	402	Total	C	N	O	S	0	2	0
			3224	2053	568	587	16			
1	B	403	Total	C	N	O	S	0	3	0
			3231	2057	570	588	16			
1	C	402	Total	C	N	O	S	0	2	0
			3220	2050	567	587	16			
1	D	403	Total	C	N	O	S	0	2	0
			3226	2054	569	587	16			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	298	GLU	ASP	variant	UNP P29474
B	298	GLU	ASP	variant	UNP P29474
C	298	GLU	ASP	variant	UNP P29474
D	298	GLU	ASP	variant	UNP P29474

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (CCD ID: HEM) (formula:  $C_{34}H_{32}FeN_4O_4$ ).




- Molecule 3 is 5,6,7,8-TETRAHYDROBIOPTERIN (CCD ID: H4B) (formula:  $\text{C}_9\text{H}_{15}\text{N}_5\text{O}_3$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total 17	C 9	N 5	O 3	0	0
3	B	1	Total 17	C 9	N 5	O 3	0	0
3	C	1	Total 17	C 9	N 5	O 3	0	0
3	D	1	Total 17	C 9	N 5	O 3	0	0

- Molecule 4 is 7-[3-(aminomethyl)-4-(cyclobutylmethoxy)phenyl]-4-methylquinolin-2-amine (CCD ID: OUP) (formula:  $C_{22}H_{25}N_3O$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total 26	C 22	N 3	O 1	0	0
4	B	1	Total 26	C 22	N 3	O 1	0	0
4	C	1	Total 26	C 22	N 3	O 1	0	0
4	D	1	Total 26	C 22	N 3	O 1	0	0

- Molecule 5 is 2-[BIS-(2-HYDROXY-ETHYL)-AMINO]-2-HYDROXYMETHYL-PROPAN E-1,3-DIOL (CCD ID: BTB) (formula:  $C_8H_{19}NO_5$ ).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	A	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	B	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	C	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		
5	D	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 6 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	Zn	0	0
			1	1		

*Continued on next page...*

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	C	1	Total	Zn	0	0
			1	1		

- Molecule 7 is GLYCEROL (CCD ID: GOL) (formula:  $C_3H_8O_3$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	A	1	Total	C	O	0	0
			6	3	3		
7	C	1	Total	C	O	0	0
			6	3	3		

- Molecule 8 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	1	Total	Cl	0	0
			1	1		
8	B	1	Total	Cl	0	0
			1	1		
8	C	1	Total	Cl	0	0
			1	1		
8	D	1	Total	Cl	0	0
			1	1		

- Molecule 9 is GADOLINIUM ATOM (CCD ID: GD) (formula: Gd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	1	Total 1	Gd 1	0	0
9	B	1	Total 1	Gd 1	0	0
9	C	1	Total 1	Gd 1	0	0
9	D	1	Total 1	Gd 1	0	0


- Molecule 10 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	A	80	Total 80	O 80	0	0
10	B	137	Total 137	O 137	0	0
10	C	111	Total 111	O 111	0	0
10	D	148	Total 148	O 148	0	0






● Molecule 1: Nitric oxide synthase, endothelial



● Molecule 1: Nitric oxide synthase, endothelial



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.07Å 152.95Å 109.23Å 90.00° 90.89° 90.00°	Depositor
Resolution (Å)	39.79 – 1.97 39.79 – 1.97	Depositor EDS
% Data completeness (in resolution range)	88.5 (39.79-1.97) 88.9 (39.79-1.97)	Depositor EDS
$R_{merge}$	0.18	Depositor
$R_{sym}$	0.18	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.18 (at 1.97Å)	Xtriage
Refinement program	PHENIX (1.11.1-2575_1496: ???)	Depositor
R, $R_{free}$	0.207 , 0.261 0.202 , 0.256	Depositor DCC
$R_{free}$ test set	6087 reflections (3.21%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	28.8	Xtriage
Anisotropy	0.317	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 54.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.116 for h,-k,-l	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	13897	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.60% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, OUP, ZN, BTB, CL, H4B, GD, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.30	0/3322	0.53	0/4524
1	B	0.34	0/3332	0.55	0/4539
1	C	0.34	1/3318 (0.0%)	0.52	0/4521
1	D	0.36	0/3324	0.56	0/4528
All	All	0.34	1/13296 (0.0%)	0.54	0/18112

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	372	ARG	CA-C	-5.14	1.50	1.53

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3224	0	3131	54	0
1	B	3231	0	3140	33	0
1	C	3220	0	3122	36	0
1	D	3226	0	3134	32	0
2	A	43	0	30	4	0
2	B	43	0	30	4	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	C	43	0	30	3	0
2	D	43	0	30	4	0
3	A	17	0	15	1	0
3	B	17	0	15	3	0
3	C	17	0	15	0	0
3	D	17	0	15	1	0
4	A	26	0	0	2	0
4	B	26	0	0	2	0
4	C	26	0	0	3	0
4	D	26	0	0	2	0
5	A	42	0	56	5	0
5	B	42	0	54	11	0
5	C	42	0	56	10	0
5	D	28	0	35	5	0
6	A	1	0	0	0	0
6	C	1	0	0	0	0
7	A	6	0	8	1	0
7	C	6	0	8	1	0
8	A	1	0	0	0	0
8	B	1	0	0	0	0
8	C	1	0	0	0	0
8	D	1	0	0	0	0
9	A	1	0	0	0	0
9	B	1	0	0	0	0
9	C	1	0	0	0	0
9	D	1	0	0	0	0
10	A	80	0	0	2	0
10	B	137	0	0	3	0
10	C	111	0	0	2	0
10	D	148	0	0	2	0
All	All	13897	0	12924	182	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 182 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:257:GLN:HA	1:B:259:GLY:H	1.31	0.95
1:C:183:ARG:HB2	2:C:501:HEM:HBD2	1.58	0.84
5:B:505:BTB:O8	10:B:601:HOH:O	2.01	0.78

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:105:PHE:O	1:D:107:ARG:N	2.22	0.72
1:B:100:LEU:HB3	1:B:103:LEU:HD22	1.70	0.71

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	400/440 (91%)	380 (95%)	14 (4%)	6 (2%)	8	2
1	B	402/440 (91%)	391 (97%)	11 (3%)	0	100	100
1	C	400/440 (91%)	387 (97%)	12 (3%)	1 (0%)	36	27
1	D	401/440 (91%)	390 (97%)	9 (2%)	2 (0%)	24	14
All	All	1603/1760 (91%)	1548 (97%)	46 (3%)	9 (1%)	21	12

5 of 9 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	90	GLN
1	A	144	GLN
1	C	89	GLN
1	A	143	SER
1	A	239	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	344/373 (92%)	331 (96%)	13 (4%)	29	19
1	B	345/373 (92%)	331 (96%)	14 (4%)	27	16
1	C	343/373 (92%)	331 (96%)	12 (4%)	32	21
1	D	344/373 (92%)	332 (96%)	12 (4%)	32	21
All	All	1376/1492 (92%)	1325 (96%)	51 (4%)	30	20

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	98	ARG
1	C	257	GLN
1	D	388	ARG
1	C	141	SER
1	C	200	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 11 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	277	HIS
1	C	283	ASN
1	D	256	GLN
1	C	408	HIS
1	C	194	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 35 ligands modelled in this entry, 10 are monoatomic - leaving 25 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
5	BTB	B	505	-	13,13,13	0.43	0	7,16,16	0.95	0
5	BTB	C	504	9	13,13,13	0.49	0	7,16,16	1.46	1 (14%)
2	HEM	A	501	1	50,50,50	1.71	9 (18%)	67,82,82	1.42	9 (13%)
5	BTB	A	506	-	13,13,13	0.46	0	7,16,16	0.36	0
3	H4B	A	502	-	17,18,18	0.79	0	14,26,26	1.90	4 (28%)
7	GOL	C	508	-	5,5,5	0.43	0	5,5,5	1.00	0
2	HEM	C	501	1	50,50,50	1.87	9 (18%)	67,82,82	1.43	9 (13%)
5	BTB	C	506	-	13,13,13	0.49	0	7,16,16	0.35	0
5	BTB	D	504	9	13,13,13	0.40	0	7,16,16	0.64	0
5	BTB	D	505	-	13,13,13	0.48	0	7,16,16	1.02	1 (14%)
2	HEM	D	501	1	50,50,50	1.76	8 (16%)	67,82,82	1.49	12 (17%)
5	BTB	A	505	-	13,13,13	0.50	0	7,16,16	0.80	0
3	H4B	C	502	-	17,18,18	0.96	0	14,26,26	1.79	4 (28%)
5	BTB	B	504	9	13,13,13	0.42	0	7,16,16	0.74	0
2	HEM	B	501	1	50,50,50	1.67	9 (18%)	67,82,82	1.49	11 (16%)
3	H4B	D	502	-	17,18,18	0.80	0	14,26,26	2.18	9 (64%)
4	OUP	B	503	-	29,29,29	0.86	0	40,41,41	1.12	2 (5%)
4	OUP	C	503	-	29,29,29	0.86	0	40,41,41	1.22	2 (5%)
5	BTB	C	505	-	13,13,13	1.01	1 (7%)	7,16,16	1.77	2 (28%)
3	H4B	B	502	-	17,18,18	0.82	0	14,26,26	2.18	6 (42%)
5	BTB	B	506	-	13,13,13	0.39	0	7,16,16	0.57	0
5	BTB	A	504	9	13,13,13	0.55	0	7,16,16	1.41	2 (28%)
4	OUP	A	503	-	29,29,29	0.84	1 (3%)	40,41,41	1.26	7 (17%)
4	OUP	D	503	-	29,29,29	0.83	0	40,41,41	1.26	4 (10%)
7	GOL	A	508	-	5,5,5	0.41	0	5,5,5	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the



Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.  
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	BTB	B	505	-	-	6/21/21/21	-
5	BTB	C	504	9	-	5/21/21/21	-
2	HEM	A	501	1	-	7/14/54/54	-
5	BTB	A	506	-	-	3/21/21/21	-
3	H4B	A	502	-	-	3/8/17/17	0/2/2/2
7	GOL	C	508	-	-	4/4/4/4	-
2	HEM	C	501	1	-	4/14/54/54	-
5	BTB	C	506	-	-	9/21/21/21	-
5	BTB	D	504	9	-	3/21/21/21	-
5	BTB	D	505	-	-	17/21/21/21	-
2	HEM	D	501	1	-	0/14/54/54	-
5	BTB	A	505	-	-	10/21/21/21	-
3	H4B	C	502	-	-	3/8/17/17	0/2/2/2
5	BTB	B	504	9	-	4/21/21/21	-
2	HEM	B	501	1	-	4/14/54/54	-
3	H4B	D	502	-	-	4/8/17/17	0/2/2/2
4	OUP	B	503	-	-	1/11/17/17	0/4/4/4
4	OUP	C	503	-	-	2/11/17/17	0/4/4/4
5	BTB	C	505	-	-	8/21/21/21	-
3	H4B	B	502	-	-	3/8/17/17	0/2/2/2
5	BTB	B	506	-	-	8/21/21/21	-
5	BTB	A	504	9	-	3/21/21/21	-
4	OUP	A	503	-	-	3/11/17/17	0/4/4/4
4	OUP	D	503	-	-	2/11/17/17	0/4/4/4
7	GOL	A	508	-	-	4/4/4/4	-

The worst 5 of 37 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	501	HEM	FE-NB	6.57	2.15	1.94
2	D	501	HEM	FE-NB	6.04	2.13	1.94
2	D	501	HEM	FE-NC	5.75	2.14	1.95
2	A	501	HEM	FE-NB	5.73	2.12	1.94
2	C	501	HEM	FE-NC	5.72	2.14	1.95

The worst 5 of 85 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	502	H4B	C2-N1-C8A	4.45	121.21	113.36
2	D	501	HEM	CBD-CAD-C3D	-4.24	100.80	112.53
3	D	502	H4B	C2-N1-C8A	4.16	120.70	113.36
4	C	503	OUP	O19-C14-C15	4.14	121.23	115.89
3	A	502	H4B	C2-N1-C8A	4.06	120.52	113.36

There are no chirality outliers.

5 of 120 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	502	H4B	C7-C6-C9-O9
3	B	502	H4B	C7-C6-C9-C10
3	C	502	H4B	C7-C6-C9-O9
3	C	502	H4B	C7-C6-C9-C10
3	D	502	H4B	N5-C6-C9-O9

There are no ring outliers.

23 monomers are involved in 58 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	B	505	BTB	4	0
5	C	504	BTB	4	0
2	A	501	HEM	4	0
5	A	506	BTB	4	0
3	A	502	H4B	1	0
7	C	508	GOL	1	0
2	C	501	HEM	3	0
5	C	506	BTB	3	0
5	D	504	BTB	1	0
5	D	505	BTB	4	0
2	D	501	HEM	4	0
5	A	505	BTB	1	0
5	B	504	BTB	5	0
2	B	501	HEM	4	0
3	D	502	H4B	1	0
4	B	503	OUP	2	0
4	C	503	OUP	3	0
5	C	505	BTB	3	0
3	B	502	H4B	3	0
5	B	506	BTB	2	0
4	A	503	OUP	2	0


*Continued on next page...*


*Continued from previous page...*


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	503	OUP	2	0
7	A	508	GOL	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.






















## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	402/440 (91%)	1.69	132 (32%) <b>1</b> <b>1</b>	26, 63, 116, 144	2 (0%)
1	B	403/440 (91%)	1.16	54 (13%) <b>7</b> <b>10</b>	21, 45, 82, 134	3 (0%)
1	C	402/440 (91%)	1.29	74 (18%) <b>3</b> <b>4</b>	25, 51, 90, 135	2 (0%)
1	D	403/440 (91%)	1.17	66 (16%) <b>4</b> <b>5</b>	28, 46, 80, 139	2 (0%)
All	All	1610/1760 (91%)	1.32	326 (20%) <b>3</b> <b>3</b>	21, 50, 99, 144	9 (0%)

The worst 5 of 326 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	120	PRO	6.9
1	A	145	ALA	5.5
1	A	346	LEU	5.5
1	D	119	ALA	4.8
1	A	163	TYR	4.8

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
5	BTB	B	506	14/14	0.52	0.17	76,100,108,112	0
5	BTB	A	506	14/14	0.56	0.19	97,113,121,121	0
5	BTB	C	506	14/14	0.58	0.18	85,109,122,123	0
5	BTB	A	505	14/14	0.67	0.20	70,89,102,102	0
7	GOL	A	508	6/6	0.68	0.14	71,83,84,87	0
5	BTB	D	505	14/14	0.72	0.15	59,79,83,84	0
5	BTB	C	505	14/14	0.77	0.17	46,64,88,90	0
5	BTB	B	505	14/14	0.78	0.16	35,74,84,93	0
7	GOL	C	508	6/6	0.80	0.14	48,73,77,83	0
4	OUP	A	503	26/26	0.82	0.18	36,71,87,99	0
3	H4B	A	502	17/17	0.82	0.15	54,63,82,84	0
5	BTB	D	504	14/14	0.85	0.16	35,66,75,78	0
3	H4B	C	502	17/17	0.85	0.16	38,58,72,76	0
3	H4B	D	502	17/17	0.86	0.14	37,60,72,77	0
3	H4B	B	502	17/17	0.86	0.14	41,58,74,77	0
4	OUP	C	503	26/26	0.87	0.15	38,56,66,68	0
8	CL	C	509	1/1	0.87	0.12	52,52,52,52	0
4	OUP	D	503	26/26	0.88	0.15	29,54,72,72	0
5	BTB	A	504	14/14	0.88	0.14	34,72,85,93	0
5	BTB	B	504	14/14	0.88	0.15	32,60,86,98	0
4	OUP	B	503	26/26	0.89	0.13	30,52,72,78	0
8	CL	B	507	1/1	0.90	0.10	52,52,52,52	0
5	BTB	C	504	14/14	0.90	0.12	35,63,81,90	0
8	CL	A	509	1/1	0.92	0.09	53,53,53,53	0
8	CL	D	506	1/1	0.92	0.10	55,55,55,55	0
2	HEM	A	501	43/43	0.93	0.13	37,58,88,97	0
2	HEM	C	501	43/43	0.95	0.12	30,44,70,81	0
2	HEM	D	501	43/43	0.95	0.10	22,35,56,63	0
2	HEM	B	501	43/43	0.95	0.11	28,37,60,70	0
6	ZN	A	507	1/1	0.98	0.03	45,45,45,45	0
9	GD	A	510	1/1	0.98	0.04	77,77,77,77	0
9	GD	C	510	1/1	0.98	0.04	79,79,79,79	0
9	GD	D	507	1/1	0.98	0.04	50,50,50,50	0
6	ZN	C	507	1/1	0.99	0.02	40,40,40,40	0
9	GD	B	508	1/1	0.99	0.03	48,48,48,48	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

**Electron density around OUP A 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OUP C 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around OUP D 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)




**Electron density around OUP B 503:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around HEM A 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





**Electron density around HEM C 501:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around HEM D 501:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)





## 6.5 Other polymers [i](#)

There are no such residues in this entry.