



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 15, 2026 – 10:16 AM UTC

PDB ID : 9PPM / pdb\_00009ppm  
Title : Mn-bound structure of the H77C variant of TriCyt2  
Authors : Eng, V.H.; Tezcan, F.A.  
Deposited on : 2025-07-21  
Resolution : 2.53 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.53 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	106	<div> <div></div> <div>71%</div> <div>28%</div> <div>.</div> </div>
1	B	106	<div> <div>74%</div> <div>25%</div> <div>.</div> </div>
1	C	106	<div> <div></div> <div>78%</div> <div>21%</div> <div>.</div> </div>
1	D	106	<div> <div>86%</div> <div>14%</div> </div>
1	E	106	<div> <div>77%</div> <div>22%</div> <div>.</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain	
1	F	106	<div> <div>%</div> <div> <div></div> <div>78%</div> <div>22%</div> </div> </div>	
1	G	106	<div> <div>3%</div> <div> <div></div> <div>77%</div> <div>23%</div> </div> </div>	
1	H	106	<div> <div>3%</div> <div> <div></div> <div>75%</div> <div>25%</div> </div> </div>	
1	I	106	<div> <div>%</div> <div> <div></div> <div>82%</div> <div>17%</div> </div> </div>	.
1	J	106	<div> <div></div> <div> <div></div> <div>68%</div> <div>31%</div> </div> </div>	.
1	K	106	<div> <div>3%</div> <div> <div></div> <div>63%</div> <div>35%</div> </div> </div>	.
1	L	106	<div> <div>4%</div> <div> <div></div> <div>75%</div> <div>25%</div> </div> </div>	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 10047 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Soluble cytochrome b562.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	B	106	Total	C	N	O	S	0	1	0
			839	525	146	163	5			
1	C	106	Total	C	N	O	S	0	1	0
			839	525	146	163	5			
1	D	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	E	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	F	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	G	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	H	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	I	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	J	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	K	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			
1	L	106	Total	C	N	O	S	0	0	0
			833	522	145	162	4			

There are 144 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	3	TRP	LEU	conflict	UNP P0ABE7
A	41	LYS	GLN	conflict	UNP P0ABE7
A	54	ALA	ASP	conflict	UNP P0ABE7
A	59	ILE	LYS	conflict	UNP P0ABE7
A	63	VAL	HIS	conflict	UNP P0ABE7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
A	69	LEU	VAL	conflict	UNP P0ABE7
A	70	TRP	GLY	conflict	UNP P0ABE7
A	73	HIS	ASP	conflict	UNP P0ABE7
A	77	CYS	LYS	variant	UNP P0ABE7
A	98	VAL	ARG	conflict	UNP P0ABE7
A	101	HIS	TYR	conflict	UNP P0ABE7
A	102	TRP	HIS	conflict	UNP P0ABE7
B	3	TRP	LEU	conflict	UNP P0ABE7
B	41	LYS	GLN	conflict	UNP P0ABE7
B	54	ALA	ASP	conflict	UNP P0ABE7
B	59	ILE	LYS	conflict	UNP P0ABE7
B	63	VAL	HIS	conflict	UNP P0ABE7
B	69	LEU	VAL	conflict	UNP P0ABE7
B	70	TRP	GLY	conflict	UNP P0ABE7
B	73	HIS	ASP	conflict	UNP P0ABE7
B	77	CYS	LYS	variant	UNP P0ABE7
B	98	VAL	ARG	conflict	UNP P0ABE7
B	101	HIS	TYR	conflict	UNP P0ABE7
B	102	TRP	HIS	conflict	UNP P0ABE7
C	3	TRP	LEU	conflict	UNP P0ABE7
C	41	LYS	GLN	conflict	UNP P0ABE7
C	54	ALA	ASP	conflict	UNP P0ABE7
C	59	ILE	LYS	conflict	UNP P0ABE7
C	63	VAL	HIS	conflict	UNP P0ABE7
C	69	LEU	VAL	conflict	UNP P0ABE7
C	70	TRP	GLY	conflict	UNP P0ABE7
C	73	HIS	ASP	conflict	UNP P0ABE7
C	77	CYS	LYS	variant	UNP P0ABE7
C	98	VAL	ARG	conflict	UNP P0ABE7
C	101	HIS	TYR	conflict	UNP P0ABE7
C	102	TRP	HIS	conflict	UNP P0ABE7
D	3	TRP	LEU	conflict	UNP P0ABE7
D	41	LYS	GLN	conflict	UNP P0ABE7
D	54	ALA	ASP	conflict	UNP P0ABE7
D	59	ILE	LYS	conflict	UNP P0ABE7
D	63	VAL	HIS	conflict	UNP P0ABE7
D	69	LEU	VAL	conflict	UNP P0ABE7
D	70	TRP	GLY	conflict	UNP P0ABE7
D	73	HIS	ASP	conflict	UNP P0ABE7
D	77	CYS	LYS	variant	UNP P0ABE7
D	98	VAL	ARG	conflict	UNP P0ABE7
D	101	HIS	TYR	conflict	UNP P0ABE7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	102	TRP	HIS	conflict	UNP P0ABE7
E	3	TRP	LEU	conflict	UNP P0ABE7
E	41	LYS	GLN	conflict	UNP P0ABE7
E	54	ALA	ASP	conflict	UNP P0ABE7
E	59	ILE	LYS	conflict	UNP P0ABE7
E	63	VAL	HIS	conflict	UNP P0ABE7
E	69	LEU	VAL	conflict	UNP P0ABE7
E	70	TRP	GLY	conflict	UNP P0ABE7
E	73	HIS	ASP	conflict	UNP P0ABE7
E	77	CYS	LYS	variant	UNP P0ABE7
E	98	VAL	ARG	conflict	UNP P0ABE7
E	101	HIS	TYR	conflict	UNP P0ABE7
E	102	TRP	HIS	conflict	UNP P0ABE7
F	3	TRP	LEU	conflict	UNP P0ABE7
F	41	LYS	GLN	conflict	UNP P0ABE7
F	54	ALA	ASP	conflict	UNP P0ABE7
F	59	ILE	LYS	conflict	UNP P0ABE7
F	63	VAL	HIS	conflict	UNP P0ABE7
F	69	LEU	VAL	conflict	UNP P0ABE7
F	70	TRP	GLY	conflict	UNP P0ABE7
F	73	HIS	ASP	conflict	UNP P0ABE7
F	77	CYS	LYS	variant	UNP P0ABE7
F	98	VAL	ARG	conflict	UNP P0ABE7
F	101	HIS	TYR	conflict	UNP P0ABE7
F	102	TRP	HIS	conflict	UNP P0ABE7
G	3	TRP	LEU	conflict	UNP P0ABE7
G	41	LYS	GLN	conflict	UNP P0ABE7
G	54	ALA	ASP	conflict	UNP P0ABE7
G	59	ILE	LYS	conflict	UNP P0ABE7
G	63	VAL	HIS	conflict	UNP P0ABE7
G	69	LEU	VAL	conflict	UNP P0ABE7
G	70	TRP	GLY	conflict	UNP P0ABE7
G	73	HIS	ASP	conflict	UNP P0ABE7
G	77	CYS	LYS	variant	UNP P0ABE7
G	98	VAL	ARG	conflict	UNP P0ABE7
G	101	HIS	TYR	conflict	UNP P0ABE7
G	102	TRP	HIS	conflict	UNP P0ABE7
H	3	TRP	LEU	conflict	UNP P0ABE7
H	41	LYS	GLN	conflict	UNP P0ABE7
H	54	ALA	ASP	conflict	UNP P0ABE7
H	59	ILE	LYS	conflict	UNP P0ABE7
H	63	VAL	HIS	conflict	UNP P0ABE7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
H	69	LEU	VAL	conflict	UNP P0ABE7
H	70	TRP	GLY	conflict	UNP P0ABE7
H	73	HIS	ASP	conflict	UNP P0ABE7
H	77	CYS	LYS	variant	UNP P0ABE7
H	98	VAL	ARG	conflict	UNP P0ABE7
H	101	HIS	TYR	conflict	UNP P0ABE7
H	102	TRP	HIS	conflict	UNP P0ABE7
I	3	TRP	LEU	conflict	UNP P0ABE7
I	41	LYS	GLN	conflict	UNP P0ABE7
I	54	ALA	ASP	conflict	UNP P0ABE7
I	59	ILE	LYS	conflict	UNP P0ABE7
I	63	VAL	HIS	conflict	UNP P0ABE7
I	69	LEU	VAL	conflict	UNP P0ABE7
I	70	TRP	GLY	conflict	UNP P0ABE7
I	73	HIS	ASP	conflict	UNP P0ABE7
I	77	CYS	LYS	variant	UNP P0ABE7
I	98	VAL	ARG	conflict	UNP P0ABE7
I	101	HIS	TYR	conflict	UNP P0ABE7
I	102	TRP	HIS	conflict	UNP P0ABE7
J	3	TRP	LEU	conflict	UNP P0ABE7
J	41	LYS	GLN	conflict	UNP P0ABE7
J	54	ALA	ASP	conflict	UNP P0ABE7
J	59	ILE	LYS	conflict	UNP P0ABE7
J	63	VAL	HIS	conflict	UNP P0ABE7
J	69	LEU	VAL	conflict	UNP P0ABE7
J	70	TRP	GLY	conflict	UNP P0ABE7
J	73	HIS	ASP	conflict	UNP P0ABE7
J	77	CYS	LYS	variant	UNP P0ABE7
J	98	VAL	ARG	conflict	UNP P0ABE7
J	101	HIS	TYR	conflict	UNP P0ABE7
J	102	TRP	HIS	conflict	UNP P0ABE7
K	3	TRP	LEU	conflict	UNP P0ABE7
K	41	LYS	GLN	conflict	UNP P0ABE7
K	54	ALA	ASP	conflict	UNP P0ABE7
K	59	ILE	LYS	conflict	UNP P0ABE7
K	63	VAL	HIS	conflict	UNP P0ABE7
K	69	LEU	VAL	conflict	UNP P0ABE7
K	70	TRP	GLY	conflict	UNP P0ABE7
K	73	HIS	ASP	conflict	UNP P0ABE7
K	77	CYS	LYS	variant	UNP P0ABE7
K	98	VAL	ARG	conflict	UNP P0ABE7
K	101	HIS	TYR	conflict	UNP P0ABE7

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
K	102	TRP	HIS	conflict	UNP P0ABE7
L	3	TRP	LEU	conflict	UNP P0ABE7
L	41	LYS	GLN	conflict	UNP P0ABE7
L	54	ALA	ASP	conflict	UNP P0ABE7
L	59	ILE	LYS	conflict	UNP P0ABE7
L	63	VAL	HIS	conflict	UNP P0ABE7
L	69	LEU	VAL	conflict	UNP P0ABE7
L	70	TRP	GLY	conflict	UNP P0ABE7
L	73	HIS	ASP	conflict	UNP P0ABE7
L	77	CYS	LYS	variant	UNP P0ABE7
L	98	VAL	ARG	conflict	UNP P0ABE7
L	101	HIS	TYR	conflict	UNP P0ABE7
L	102	TRP	HIS	conflict	UNP P0ABE7

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total Na 1 1	0	0

- Molecule 3 is MANGANESE (II) ION (CCD ID: MN) (formula: Mn) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	C	1	Total Mn 1 1	0	0
3	E	1	Total Mn 1 1	0	0
3	F	1	Total Mn 1 1	0	0
3	J	1	Total Mn 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	4	Total O 4 4	0	0
4	B	2	Total O 2 2	0	0
4	C	6	Total O 6 6	0	0

*Continued on next page...*




*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	D	3	Total 3	O 3	0	0
4	E	4	Total 4	O 4	0	0
4	F	1	Total 1	O 1	0	0
4	G	2	Total 2	O 2	0	0
4	H	1	Total 1	O 1	0	0
4	I	5	Total 5	O 5	0	0
4	J	3	Total 3	O 3	0	0
4	K	1	Total 1	O 1	0	0
4	L	2	Total 2	O 2	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562





- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562




- Molecule 1: Soluble cytochrome b562



- Molecule 1: Soluble cytochrome b562



- Molecule 1: Soluble cytochrome b562





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	71.19Å 47.36Å 166.23Å 90.00° 97.27° 90.00°	Depositor
Resolution (Å)	46.31 – 2.53 46.31 – 2.53	Depositor EDS
% Data completeness (in resolution range)	99.2 (46.31-2.53) 99.5 (46.31-2.53)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.72 (at 2.51Å)	Xtriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.234 , 0.291 0.234 , 0.291	Depositor DCC
$R_{free}$ test set	1999 reflections (5.35%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.0	Xtriage
Anisotropy	0.108	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 40.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	10047	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 9.72% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.18	0/847	0.37	0/1144
1	B	0.14	0/853	0.28	0/1152
1	C	0.21	0/853	0.36	0/1152
1	D	0.13	0/847	0.26	0/1144
1	E	0.13	0/847	0.27	0/1144
1	F	0.12	0/847	0.26	0/1144
1	G	0.16	0/847	0.33	0/1144
1	H	0.15	0/847	0.31	0/1144
1	I	0.14	0/847	0.33	0/1144
1	J	0.15	0/847	0.33	0/1144
1	K	0.14	0/847	0.33	0/1144
1	L	0.17	0/847	0.38	0/1144
All	All	0.15	0/10176	0.32	0/13744

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	833	0	832	21	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	839	0	836	21	0
1	C	839	0	836	15	0
1	D	833	0	832	10	0
1	E	833	0	832	18	0
1	F	833	0	832	15	0
1	G	833	0	832	19	0
1	H	833	0	832	20	0
1	I	833	0	832	12	0
1	J	833	0	832	26	0
1	K	833	0	832	26	0
1	L	833	0	832	22	0
2	A	1	0	0	0	0
3	C	1	0	0	0	0
3	E	1	0	0	0	0
3	F	1	0	0	0	0
3	J	1	0	0	0	0
4	A	4	0	0	1	0
4	B	2	0	0	0	0
4	C	6	0	0	0	0
4	D	3	0	0	0	0
4	E	4	0	0	1	0
4	F	1	0	0	0	0
4	G	2	0	0	0	0
4	H	1	0	0	0	0
4	I	5	0	0	0	0
4	J	3	0	0	0	0
4	K	1	0	0	0	0
4	L	2	0	0	0	0
All	All	10047	0	9992	206	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 206 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:TRP:HE1	1:B:101:HIS:HE1	1.23	0.87
1:K:83:LYS:HG3	1:K:86:GLU:HB2	1.57	0.85
1:F:63:VAL:HG21	1:K:42:LYS:HD2	1.70	0.73
1:J:22:ASN:H	1:J:25:GLN:HE21	1.36	0.71
1:J:17:ILE:HG12	1:J:26:VAL:HG13	1.74	0.70

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	104/106 (98%)	102 (98%)	1 (1%)	1 (1%)	12	23
1	B	105/106 (99%)	102 (97%)	3 (3%)	0	100	100
1	C	105/106 (99%)	103 (98%)	2 (2%)	0	100	100
1	D	104/106 (98%)	104 (100%)	0	0	100	100
1	E	104/106 (98%)	101 (97%)	3 (3%)	0	100	100
1	F	104/106 (98%)	103 (99%)	1 (1%)	0	100	100
1	G	104/106 (98%)	101 (97%)	3 (3%)	0	100	100
1	H	104/106 (98%)	103 (99%)	1 (1%)	0	100	100
1	I	104/106 (98%)	103 (99%)	1 (1%)	0	100	100
1	J	104/106 (98%)	95 (91%)	9 (9%)	0	100	100
1	K	104/106 (98%)	98 (94%)	6 (6%)	0	100	100
1	L	104/106 (98%)	100 (96%)	4 (4%)	0	100	100
All	All	1250/1272 (98%)	1215 (97%)	34 (3%)	1 (0%)	48	67

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	100	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/86 (100%)	84 (98%)	2 (2%)	44	70
1	B	87/86 (101%)	86 (99%)	1 (1%)	65	83
1	C	87/86 (101%)	84 (97%)	3 (3%)	32	57
1	D	86/86 (100%)	85 (99%)	1 (1%)	63	82
1	E	86/86 (100%)	84 (98%)	2 (2%)	44	70
1	F	86/86 (100%)	85 (99%)	1 (1%)	63	82
1	G	86/86 (100%)	86 (100%)	0	100	100
1	H	86/86 (100%)	85 (99%)	1 (1%)	63	82
1	I	86/86 (100%)	84 (98%)	2 (2%)	44	70
1	J	86/86 (100%)	85 (99%)	1 (1%)	63	82
1	K	86/86 (100%)	81 (94%)	5 (6%)	18	36
1	L	86/86 (100%)	85 (99%)	1 (1%)	63	82
All	All	1034/1032 (100%)	1014 (98%)	20 (2%)	51	74

5 of 20 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	K	57	GLU
1	K	88	GLN
1	L	60	ASP
1	K	95	LYS
1	D	77	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	11	ASN
1	J	25	GLN
1	L	88	GLN
1	J	88	GLN
1	I	103	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 5 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	106/106 (100%)	0.20	1 (0%) 81 79	39, 58, 76, 91	0
1	B	106/106 (100%)	0.08	0 100 100	27, 52, 74, 82	1 (0%)
1	C	106/106 (100%)	0.13	1 (0%) 81 79	28, 57, 75, 93	1 (0%)
1	D	106/106 (100%)	0.13	0 100 100	39, 56, 72, 88	0
1	E	106/106 (100%)	0.19	0 100 100	39, 59, 76, 88	0
1	F	106/106 (100%)	0.12	1 (0%) 81 79	38, 58, 76, 88	0
1	G	106/106 (100%)	0.41	3 (2%) 55 51	40, 67, 89, 95	0
1	H	106/106 (100%)	0.41	3 (2%) 55 51	44, 69, 89, 102	0
1	I	106/106 (100%)	0.49	1 (0%) 81 79	50, 70, 97, 106	0
1	J	106/106 (100%)	0.46	0 100 100	49, 75, 95, 105	0
1	K	106/106 (100%)	0.74	3 (2%) 55 51	50, 89, 106, 119	0
1	L	106/106 (100%)	0.58	4 (3%) 44 40	49, 72, 96, 113	0
All	All	1272/1272 (100%)	0.33	17 (1%) 75 72	27, 64, 95, 119	2 (0%)

The worst 5 of 17 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	87	ALA	3.3
1	L	1	ALA	3.1
1	G	1	ALA	2.9
1	C	2	ASP	2.7
1	A	1	ALA	2.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

### 6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
3	MN	J	201	1/1	0.88	0.20	76,76,76,76	1
2	NA	A	201	1/1	0.89	0.28	65,65,65,65	0
3	MN	F	201	1/1	0.90	0.21	72,72,72,72	1
3	MN	C	201	1/1	0.92	0.18	66,66,66,66	0
3	MN	E	201	1/1	0.93	0.18	68,68,68,68	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




**Electron density around MN J 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MN F 201:**




$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)






**Electron density around MN C 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



**Electron density around MN E 201:**

$2mF_o-DF_c$  (at 0.7 rmsd) in gray  
 $mF_o-DF_c$  (at 3 rmsd) in purple (negative)  
and green (positive)



## 6.5 Other polymers ⓘ

There are no such residues in this entry.