



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:11 AM UTC

PDB ID : 6TKF / pdb_00006tkf
Title : ChiLob 7/4 H2 HC-C225S KappaLC-C214S F(ab')₂
Authors : Orr, C.M.; Fisher, H.; Tews, I.
Deposited on : 2019-11-28
Resolution : 2.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	8975 (2.20-2.16)
Clashscore	190562	9786 (2.20-2.16)
Ramachandran outliers	187476	9664 (2.20-2.16)
Sidechain outliers	187428	9664 (2.20-2.16)
RSRZ outliers	180081	8979 (2.20-2.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	HHH	231	
2	LLL	214	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6716 atoms, of which 3190 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ChiLob 7/4 H2 Heavy chain C225S.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	HHH	216	Total	C	H	N	O	S	106	2	0
			3242	1032	1604	270	325	11			

- Molecule 2 is a protein called ChiLob 7/4 H2 Kappa light chain C214S.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	LLL	212	Total	C	H	N	O	S	119	1	0
			3226	1025	1586	272	338	5			

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	HHH	75	Total	O	0	0
			75	75		
3	LLL	173	Total	O	0	0
			173	173		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ChiLob 7/4 H2 Heavy chain C225S

Chain HHH: 



- Molecule 2: ChiLob 7/4 H2 Kappa light chain C214S

Chain LLL: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants a, b, c, α , β , γ	149.10Å 149.10Å 45.80Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.85 – 2.18 48.85 – 2.18	Depositor EDS
% Data completeness (in resolution range)	99.8 (48.85-2.18) 99.8 (48.85-2.18)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.10 (at 2.18Å)	Xtriage
Refinement program	REFMAC 5.8.0258	Depositor
R, R_{free}	0.196 , 0.239 0.196 , 0.239	Depositor DCC
R_{free} test set	1490 reflections (4.58%)	wwPDB-VP
Wilson B-factor (Å ²)	35.7	Xtriage
Anisotropy	0.247	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.029 for -h,-k,l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6716	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.43% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	HHH	1.04	0/1681	1.31	2/2285 (0.1%)
2	LLL	1.09	3/1678 (0.2%)	1.26	4/2281 (0.2%)
All	All	1.07	3/3359 (0.1%)	1.29	6/4566 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	LLL	120	PRO	C-O	-5.48	1.17	1.23
2	LLL	32	TYR	C-O	5.47	1.29	1.23
2	LLL	132	VAL	C-O	5.09	1.29	1.24

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	HHH	49	GLY	CA-C-N	6.31	126.47	121.61
1	HHH	49	GLY	C-N-CA	6.31	126.47	121.61
2	LLL	141	PRO	CA-C-N	6.10	128.45	120.28
2	LLL	141	PRO	C-N-CA	6.10	128.45	120.28
2	LLL	93	ASN	CB-CA-C	-5.33	98.84	109.33
2	LLL	180	THR	CA-CB-OG1	-5.21	101.78	109.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	HHH	1638	1604	1589	10	0
2	LLL	1640	1586	1583	11	0
3	HHH	75	0	0	1	0
3	LLL	173	0	0	1	0
All	All	3526	3190	3172	21	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (21) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:HHH:48:ILE:HD13	1:HHH:81:MET:HE1	1.70	0.73
1:HHH:81:MET:CE	1:HHH:83:LEU:HD21	2.26	0.66
2:LLL:40:PRO:HB3	2:LLL:165:GLU:HG3	1.78	0.63
1:HHH:135:PRO:O	1:HHH:223:LYS:HE2	1.99	0.62
1:HHH:81:MET:HE2	1:HHH:83:LEU:HD21	1.92	0.51
1:HHH:187:LEU:C	1:HHH:187:LEU:HD12	2.37	0.50
2:LLL:4:MET:HE3	2:LLL:23:CYS:SG	2.52	0.50
2:LLL:186:TYR:CZ	2:LLL:211:ARG:HD3	2.47	0.49
2:LLL:12:SER:HA	2:LLL:105:GLU:O	2.13	0.49
1:HHH:201:GLN:HB2	3:HHH:357:HOH:O	2.13	0.49
2:LLL:24:SER:HA	2:LLL:69:THR:O	2.14	0.47
2:LLL:186:TYR:CE2	2:LLL:211:ARG:HD3	2.50	0.46
2:LLL:33:LEU:C	2:LLL:33:LEU:HD13	2.39	0.46
2:LLL:186:TYR:CE2	2:LLL:211:ARG:CD	3.01	0.43
2:LLL:33:LEU:HD21	2:LLL:88:CYS:HB2	2.00	0.43
2:LLL:54:LEU:HD21	2:LLL:60:SER:HA	2.01	0.43
1:HHH:205[A]:CYS:SG	1:HHH:218:LYS:HB3	2.59	0.42
2:LLL:38:GLN:NE2	3:LLL:308:HOH:O	2.52	0.42
1:HHH:48:ILE:HG21	1:HHH:81:MET:HE1	2.01	0.42
1:HHH:201:GLN:O	1:HHH:222:ARG:NH2	2.53	0.41
1:HHH:81:MET:HE3	1:HHH:83:LEU:HD21	2.02	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	HHH	212/231 (92%)	205 (97%)	7 (3%)	0	100	100
2	LLL	211/214 (99%)	205 (97%)	6 (3%)	0	100	100
All	All	423/445 (95%)	410 (97%)	13 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	HHH	187/203 (92%)	183 (98%)	4 (2%)	47	59
2	LLL	189/190 (100%)	188 (100%)	1 (0%)	81	89
All	All	376/393 (96%)	371 (99%)	5 (1%)	61	74

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	HHH	17	SER
1	HHH	85	SER
1	HHH	143	SER
1	HHH	206	ASN
2	LLL	83	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	HHH	216/231 (93%)	0.38	9 (4%) 40 40	17, 47, 70, 80	2 (0%)
2	LLL	212/214 (99%)	-0.49	1 (0%) 87 86	15, 29, 47, 68	1 (0%)
All	All	428/445 (96%)	-0.05	10 (2%) 61 60	15, 35, 65, 80	3 (0%)

All (10) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	HHH	106	TYR	4.2
1	HHH	142	GLU	3.7
1	HHH	101	VAL	3.6
1	HHH	136	CYS	3.5
1	HHH	107	TYR	3.2
1	HHH	102	TYR	3.2
2	LLL	212	GLY	3.0
1	HHH	224	CYS	2.9
1	HHH	200	THR	2.8
1	HHH	196	SER	2.4

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.