



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 10:43 PM UTC

PDB ID : 6UTV / pdb\_00006utv  
Title : E. coli sigma-S transcription initiation complex with a 6-nt RNA ("Fresh" crystal soaked with CTP, UTP, GTP, and ddATP for 150 seconds)  
Authors : Zuo, Y.; De, S.; Steitz, T.A.  
Deposited on : 2019-10-30  
Resolution : 3.45 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.








Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1070 (3.50-3.42)
Clashscore	190562	1128 (3.50-3.42)
Ramachandran outliers	187476	1101 (3.50-3.42)
Sidechain outliers	187428	1102 (3.50-3.42)
RSRZ outliers	180081	1070 (3.50-3.42)
RNA backbone	3983	1186 (3.92-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	AAA	242	 69% 23% • 5%
1	BBB	242	 56% 34% • 6%
2	CCC	1342	 81% 18% •
3	DDD	1407	 76% 19% •

Continued on next page...

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
4	EEE	90	 76% 11% 12%
5	FFF	336	 60% 22% 18%
6	111	50	 26% 34% 40%
7	222	50	 38% 32% 30%
8	333	6	 50% 33% 17%

## 2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 29063 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	AAA	230	Total	C	N	O	S	0	0	0
			1787	1112	317	352	6			
1	BBB	228	Total	C	N	O	S	0	0	0
			1767	1100	312	349	6			

There are 14 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
AAA	-6	ALA	-	expression tag	UNP A0A377D9Q8
AAA	-5	HIS	-	expression tag	UNP A0A377D9Q8
AAA	-4	HIS	-	expression tag	UNP A0A377D9Q8
AAA	-3	HIS	-	expression tag	UNP A0A377D9Q8
AAA	-2	HIS	-	expression tag	UNP A0A377D9Q8
AAA	-1	HIS	-	expression tag	UNP A0A377D9Q8
AAA	0	HIS	-	expression tag	UNP A0A377D9Q8
BBB	-6	ALA	-	expression tag	UNP A0A377D9Q8
BBB	-5	HIS	-	expression tag	UNP A0A377D9Q8
BBB	-4	HIS	-	expression tag	UNP A0A377D9Q8
BBB	-3	HIS	-	expression tag	UNP A0A377D9Q8
BBB	-2	HIS	-	expression tag	UNP A0A377D9Q8
BBB	-1	HIS	-	expression tag	UNP A0A377D9Q8
BBB	0	HIS	-	expression tag	UNP A0A377D9Q8

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	CCC	1341	Total	C	N	O	S	0	0	0
			10577	6636	1842	2056	43			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	DDD	1362	Total	C	N	O	S	0	0	0
			10568	6633	1887	1998	50			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	EEE	79	Total	C	N	O	S	0	0	0
			627	382	118	126	1			

- Molecule 5 is a protein called RNA polymerase sigma factor RpoS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	FFF	277	Total	C	N	O	S	0	0	0
			2253	1411	415	423	4			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
FFF	2	GLY	SER	conflict	UNP P13445
FFF	33	GLU	GLN	conflict	UNP P13445
FFF	329	LEU	-	expression tag	UNP P13445
FFF	330	GLU	-	expression tag	UNP P13445
FFF	331	HIS	-	expression tag	UNP P13445
FFF	332	HIS	-	expression tag	UNP P13445
FFF	333	HIS	-	expression tag	UNP P13445
FFF	334	HIS	-	expression tag	UNP P13445
FFF	335	HIS	-	expression tag	UNP P13445
FFF	336	HIS	-	expression tag	UNP P13445

- Molecule 6 is a DNA chain called Synthetic DNA 50-MER (promoter non-template strand).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	111	30	Total	C	N	O	P	0	0	0
			618	294	111	183	30			

- Molecule 7 is a DNA chain called Synthetic DNA 50-MER (Promoter template strand).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	222	35	Total	C	N	O	P	0	0	0
			716	342	132	208	34			

- Molecule 8 is a RNA chain called RNA 6-mer.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	333	6	Total	C	N	O	P	0	0	0
			137	57	22	50	8			


- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	CCC	1	Total	Mg	0	0
			1	1		
9	DDD	1	Total	Mg	0	0
			1	1		

- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	DDD	2	Total	Zn	0	0
			2	2		

- Molecule 11 is DIPHOSPHATE (CCD ID: DPO) (formula: O<sub>7</sub>P<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
11	DDD	1	Total	O	P	0	0
			9	7	2		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


#### • Molecule 1: DNA-directed RNA polymerase subunit alpha




#### • Molecule 1: DNA-directed RNA polymerase subunit alpha




#### • Molecule 2: DNA-directed RNA polymerase subunit beta





• Molecule 3: DNA-directed RNA polymerase subunit beta'


Chain DDD: 76% 19% . .





GLY  
SER  
ASP  
ASN  
GLU

- Molecule 4: DNA-directed RNA polymerase subunit omega

Chain EEE:  76% 11% 12%

A2 R3 T5 R28 E41 E42 K45 T46 T47 Q73 E76 L80 GLN ALA VAL THR ALA ILE ALA GLY ARG ARG

- Molecule 5: RNA polymerase sigma factor RpoS

Chain FFF:  60% 22% 18%


NET GLY GLN ASN THR LEU LYS VAL HIS ASP LEU ASN GLU ASP ALA PHE GLY VAL GLU VAL PHE ASP GLY LYS ALA VAL LEU VAL GLU PRO SER ASP ASN ASP LEU ALA GLU GLU LEU LEU SER LEU GLN THR R53 V54 E74 L84 R108

G113 L114 A115 L116 L117 D118 L119 L120 E121 K133 P136 S143 T147 Q152 T153 R156 M159 T162 T163 T164 T165 T166 L167 P168 I169 H170 I171 V172 K173 M176 V177 Y178 A182 L189 D190 H191 E192 P193 D203 V209 M212 L213 R214 T219

T220 S221 V222 L226 G227 G228 G229 S230 L234 L235 E241 E242 E243 P246 E247 T250 Q251 V262 A268 R271 E272 A277 F278 G279 L280 T287 V291 G292 T295 G296 L297 T298 R301 Q306 Q320 N323 T324 E325 L329 GLU HIS HIS HIS

HIS  
HIS  
HIS

- Molecule 6: Synthetic DNA 50-MER (promoter non-template strand)

Chain 111:  26% 34% 40%

DA DC DC DT DT DG DA DC DC DC DA DC DC DC T26 C29 G30 T31 A32 T33 G34 G35 T36 A37 T38 A39 A40 T41 G42 T43 G44 DT DC DC DA G49 T50 C51 T52 G53 A54 C55 G56 G59

- Molecule 7: Synthetic DNA 50-MER (Promoter template strand)

Chain 222:  38% 32% 30%

T3 G8 T9 G12 A13 C14 T15 C16 G17 T18 A19 DG DG A22 T23 T24 A25 T26 A27 G28 A32 C33 G39 DT DG DG DG DG DA DT DG DG DC DA DA DA DG

- Molecule 8: RNA 6-mer

Chain 333:  50% 33% 17%

G14 C18 U19

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	132.53Å 155.43Å 232.60Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.28 – 3.45 49.28 – 3.45	Depositor EDS
% Data completeness (in resolution range)	99.2 (49.28-3.45) 99.2 (49.28-3.45)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.06 (at 3.48Å)	Xtriage
Refinement program	REFMAC 5.8.0257	Depositor
R, $R_{free}$	0.282 , 0.348 0.277 , 0.339	Depositor DCC
$R_{free}$ test set	3029 reflections (4.74%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	121.7	Xtriage
Anisotropy	0.265	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 120.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.36$ , $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	29063	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	195.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.32% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, DPO, GTP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	AAA	0.99	0/1809	1.17	1/2450 (0.0%)
1	BBB	1.02	0/1789	1.19	1/2425 (0.0%)
2	CCC	0.93	0/10746	1.28	7/14499 (0.0%)
3	DDD	0.94	0/10729	1.28	8/14487 (0.1%)
4	EEE	0.90	0/629	1.35	0/847
5	FFF	0.99	0/2282	1.32	0/3076
6	111	0.39	0/691	0.67	0/1063
7	222	0.38	0/802	0.65	0/1234
8	333	0.46	0/116	0.88	0/178
All	All	0.93	0/29593	1.25	17/40259 (0.0%)

There are no bond length outliers.

The worst 5 of 17 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	DDD	504	GLN	CB-CA-C	-5.95	109.14	117.23
3	DDD	777	HIS	N-CA-C	-5.67	105.00	111.07
2	CCC	1261	GLY	CA-C-O	-5.60	118.36	122.23
2	CCC	181	GLY	CA-C-O	-5.53	117.21	122.13
3	DDD	462	ASP	N-CA-C	-5.41	104.95	112.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	AAA	1787	0	1813	71	0
1	BBB	1767	0	1789	144	0
2	CCC	10577	0	10591	225	0
3	DDD	10568	0	10780	262	1
4	EEE	627	0	634	14	0
5	FFF	2253	0	2298	74	0
6	111	618	0	341	28	0
7	222	716	0	397	29	1
8	333	137	0	65	5	0
9	CCC	1	0	0	0	0
9	DDD	1	0	0	0	0
10	DDD	2	0	0	0	0
11	DDD	9	0	0	1	0
All	All	29063	0	28708	691	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 691 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:DDD:817:HIS:HB3	3:DDD:860:ARG:NH2	1.16	1.40
1:BBB:49:SER:O	1:BBB:151:GLY:HA3	1.38	1.24
6:111:54:DA:H2''	6:111:55:DC:C5	1.75	1.19
3:DDD:817:HIS:CB	3:DDD:860:ARG:NH2	2.05	1.18
1:BBB:179:PRO:HG2	1:BBB:211:ILE:HD12	1.15	1.13

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:DDD:1170:LYS:NZ	7:222:33:DC:OP1[3_644]	2.16	0.04

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	AAA	228/242 (94%)	210 (92%)	12 (5%)	6 (3%)	4	27
1	BBB	226/242 (93%)	203 (90%)	20 (9%)	3 (1%)	9	39
2	CCC	1339/1342 (100%)	1245 (93%)	85 (6%)	9 (1%)	18	51
3	DDD	1360/1407 (97%)	1260 (93%)	91 (7%)	9 (1%)	18	51
4	EEE	77/90 (86%)	75 (97%)	2 (3%)	0	100	100
5	FFF	275/336 (82%)	256 (93%)	14 (5%)	5 (2%)	6	34
All	All	3505/3659 (96%)	3249 (93%)	224 (6%)	32 (1%)	14	46

5 of 32 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	AAA	192	VAL
3	DDD	519	ASN
3	DDD	1053	LEU
5	FFF	190	ASP
1	AAA	210	THR

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	AAA	198/208 (95%)	189 (96%)	9 (4%)	24	52
1	BBB	196/208 (94%)	181 (92%)	15 (8%)	12	38
2	CCC	1156/1157 (100%)	1136 (98%)	20 (2%)	53	70

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
3	DDD	1135/1168 (97%)	1116 (98%)	19 (2%)	53	70
4	EEE	67/74 (90%)	66 (98%)	1 (2%)	57	72
5	FFF	240/292 (82%)	231 (96%)	9 (4%)	29	57
All	All	2992/3107 (96%)	2919 (98%)	73 (2%)	43	65

5 of 73 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	DDD	731	ARG
5	FFF	271	ARG
3	DDD	798	ARG
5	FFF	152	GLN
2	CCC	12	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	333	4/6 (66%)	2 (50%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
8	333	18	C
8	333	19	U

There are no RNA pucker outliers to report.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry

Of 5 ligands modelled in this entry, 4 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
11	DPO	DDD	1504	-	6,8,8	0.79	0	12,13,13	0.81	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
11	DPO	DDD	1504	-	-	1/6/6/6	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
11	DDD	1504	DPO	P2-O4-P1-O1

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
11	DDD	1504	DPO	1	0

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	AAA	230/242 (95%)	-0.66	0 100 100	159, 207, 265, 284	0
1	BBB	228/242 (94%)	-0.52	3 (1%) 75 51	162, 219, 286, 326	0
2	CCC	1341/1342 (99%)	-0.81	3 (0%) 91 83	95, 172, 271, 347	0
3	DDD	1362/1407 (96%)	-0.81	1 (0%) 92 88	101, 185, 277, 345	0
4	EEE	79/90 (87%)	-0.85	0 100 100	156, 223, 347, 400	0
5	FFF	277/336 (82%)	-0.80	0 100 100	136, 210, 288, 306	0
6	111	30/50 (60%)	-1.07	0 100 100	191, 228, 301, 336	0
7	222	35/50 (70%)	-0.88	0 100 100	129, 234, 353, 421	0
8	333	5/6 (83%)	-1.45	0 100 100	137, 138, 155, 188	0
All	All	3587/3765 (95%)	-0.78	7 (0%) 91 83	95, 191, 283, 421	0

The worst 5 of 7 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	CCC	524	ILE	3.4
1	BBB	26	VAL	3.3
1	BBB	39	LEU	3.0
3	DDD	472	LEU	2.7
2	CCC	27	LEU	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
11	DPO	DDD	1504	9/9	0.90	0.04	155,178,196,201	0
9	MG	CCC	1401	1/1	0.93	0.08	156,156,156,156	0
10	ZN	DDD	1501	1/1	0.98	0.06	267,267,267,267	0
10	ZN	DDD	1502	1/1	0.99	0.04	162,162,162,162	0
9	MG	DDD	1503	1/1	1.00	0.01	93,93,93,93	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.