



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 03:42 PM UTC

PDB ID : 7VBL / pdb_00007vbl
EMDB ID : EMD-31881
Title : Membrane arm of active state CI from DQ-NADH dataset
Authors : Gu, J.K.; Yang, M.J.
Deposited on : 2021-08-31
Resolution : 2.60 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	8728 (2.10 - 3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Q	44	
2	S	70	
3	U	83	
4	V	140	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	W	113	
6	X	88	
7	Y	67	
8	Z	80	
9	a	138	
10	b	126	
11	c	156	
12	d	175	
13	e	104	
14	f	49	
15	g	122	
16	h	105	
17	i	347	
18	j	115	
19	k	98	
20	l	606	
21	m	175	
22	n	56	
23	o	128	
24	p	178	
25	r	459	
26	s	318	
27	u	171	
28	v	125	
29	w	320	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
30	PEE	l	704	-	-	X	-
31	CDL	l	702	-	-	X	-
31	CDL	r	505	-	-	X	-
33	PLX	r	503	-	-	X	-

2 Entry composition [i](#)

There are 36 unique types of molecules in this entry. The entry contains 40142 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Q	44	Total	C	N	O	S	0	0
			363	236	60	66	1		

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	S	70	Total	C	N	O	S	0	0
			566	364	103	94	5		

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	U	83	Total	C	N	O	S	0	0
			643	417	110	115	1		

- Molecule 4 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	V	140	Total	C	N	O	S	0	0
			1021	651	174	190	6		

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	W	113	Total	C	N	O	S	0	0
			949	614	160	167	8		

- Molecule 6 is a protein called Acyl carrier protein, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	X	88	Total	C	N	O	S	0	0
			691	445	103	138	5		

- Molecule 7 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	Y	67	Total	C	N	O	S	0	0
			584	385	95	103	1		

- Molecule 8 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	Z	80	Total	C	N	O	S	0	0
			611	397	107	106	1		

- Molecule 9 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	a	138	Total	C	N	O	S	0	0
			1151	754	195	199	3		

- Molecule 10 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	b	98	Total	C	N	O	S	0	0
			819	537	144	137	1		

- Molecule 11 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	c	156	Total	C	N	O	S	0	0
			1315	853	213	241	8		

- Molecule 12 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	d	175	Total	C	N	O	S	0	0
			1461	916	265	272	8		

- Molecule 13 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	e	104	Total	C	N	O	S	0	0
			867	553	142	168	4		

- Molecule 14 is a protein called NADH dehydrogenase [ubiquinone] 1 subunit C1, mitochondrial.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	f	49	Total	C	N	O	0	0
			377	246	65	66		

- Molecule 15 is a protein called NADH dehydrogenase [ubiquinone] 1 subunit C2.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	g	122	Total	C	N	O	S	0	0
			1005	653	174	172	6		

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	h	105	Total	C	N	O	S	0	0
			867	550	161	150	6		

- Molecule 17 is a protein called NADH-ubiquinone oxidoreductase chain 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	i	347	Total	C	N	O	S	0	0
			2710	1782	420	462	46		

- Molecule 18 is a protein called NADH-ubiquinone oxidoreductase chain 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	j	115	Total	C	N	O	S	0	0
			914	615	134	158	7		

- Molecule 19 is a protein called NADH-ubiquinone oxidoreductase chain 4L.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	k	98	Total	C	N	O	S	0	0
			748	493	113	128	14		

- Molecule 20 is a protein called NADH-ubiquinone oxidoreductase chain 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	l	606	Total	C	N	O	S	0	0
			4797	3181	743	822	51		

- Molecule 21 is a protein called NADH-ubiquinone oxidoreductase chain 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	m	175	Total	C	N	O	S	0	0
			1273	847	187	226	13		

- Molecule 22 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	n	56	Total	C	N	O	S	0	0
			479	311	88	79	1		

- Molecule 23 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	o	128	Total	C	N	O	S	0	0
			1062	691	182	189			

- Molecule 24 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	p	178	Total	C	N	O	S	0	0
			1534	982	279	265	8		

- Molecule 25 is a protein called NADH-ubiquinone oxidoreductase chain 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	r	459	Total	C	N	O	S	0	0
			3631	2412	572	609	38		

- Molecule 26 is a protein called NADH-ubiquinone oxidoreductase chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	s	318	Total	C	N	O	S	0	0
			2508	1678	385	424	21		

- Molecule 27 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	u	171	Total	C	N	O	S	0	0
			1398	887	250	251	10		

- Molecule 28 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	v	124	Total	C	N	O	S	0	0
			1028	642	195	182	9		


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
v	1	MYR	-	acetylation	UNP F1SCH1

- Molecule 29 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial.


Mol	Chain	Residues	Atoms					AltConf	Trace
29	w	320	Total	C	N	O	S	0	0
			2590	1649	440	491	10		

- Molecule 30 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (CCD ID: PEE) (formula: C₄₁H₇₈NO₈P) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
30	U	1	Total	C	N	O	P	0
			51	41	1	8	1	
30	U	1	Total	C	N	O	P	0
			51	41	1	8	1	
30	W	1	Total	C	N	O	P	0
			41	31	1	8	1	
30	i	1	Total	C	N	O	P	0
			47	37	1	8	1	
30	j	1	Total	C	N	O	P	0
			41	31	1	8	1	
30	l	1	Total	C	N	O	P	0
			40	30	1	8	1	
30	l	1	Total	C	N	O	P	0
			51	41	1	8	1	
30	l	1	Total	C	N	O	P	0
			46	36	1	8	1	
30	r	1	Total	C	N	O	P	0
			51	41	1	8	1	

- Molecule 31 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
31	V	1	Total	C	O	P	0
			94	75	17	2	
31	V	1	Total	C	O	P	0
			100	81	17	2	
31	a	1	Total	C	O	P	0
			100	81	17	2	
31	i	1	Total	C	O	P	0
			87	68	17	2	
31	l	1	Total	C	O	P	0
			94	75	17	2	
31	m	1	Total	C	O	P	0
			100	81	17	2	
31	r	1	Total	C	O	P	0
			99	80	17	2	
31	r	1	Total	C	O	P	0
			100	81	17	2	
31	s	1	Total	C	O	P	0
			89	70	17	2	
31	u	1	Total	C	O	P	0
			55	36	17	2	

- Molecule 32 is S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonooxy)butanoyl]-beta-alanyl}amino)ethyl] dodecanethioate (CCD ID: 8Q1) (formula: $C_{23}H_{45}N_2O_8PS$) (labeled as "Ligand of Interest" by depositor).

Continued from previous page...

Mol	Chain	Residues	Atoms					AltConf
33	g	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	j	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	j	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	r	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	r	1	Total	C	N	O	P	0
			52	42	1	8	1	

- Molecule 34 is Coenzyme Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-isomer (CCD ID: UQ) (formula: C₅₉H₉₀O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
34	s	1	Total	C	O	0
			38	34	4	

- Molecule 35 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltCon
35	w	1	Total	C	N	O	P	0
			27	10	5	10	2	

- Molecule 36 is water.

Mol	Chain	Residues	Atoms		AltConf
36	Q	5	Total 5	O 5	0
36	S	4	Total 4	O 4	0
36	U	2	Total 2	O 2	0
36	W	1	Total 1	O 1	0
36	a	1	Total 1	O 1	0
36	b	1	Total 1	O 1	0
36	c	3	Total 3	O 3	0
36	d	3	Total 3	O 3	0
36	e	1	Total 1	O 1	0
36	h	3	Total 3	O 3	0
36	i	74	Total 74	O 74	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
36	j	33	Total 33	O 33	0
36	k	35	Total 35	O 35	0
36	l	60	Total 60	O 60	0
36	m	16	Total 16	O 16	0
36	n	2	Total 2	O 2	0
36	p	2	Total 2	O 2	0
36	r	91	Total 91	O 91	0
36	s	92	Total 92	O 92	0
36	w	2	Total 2	O 2	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH dehydrogenase [ubiquinone] iron-sulfur protein 2, mitochondrial




- Molecule 2: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1




- Molecule 3: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3



- Molecule 4: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 11





- Molecule 5: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13





- Molecule 6: Acyl carrier protein, mitochondrial




- Molecule 7: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 2, mitochondrial




- Molecule 8: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 3





- Molecule 9: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 5, mitochondrial



- Molecule 10: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6



- Molecule 11: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial



E185
I186

- Molecule 12: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10

Chain d: 6% 83% 16%

P2 D3 S4 R15 Q23 T46 E50 R60 Y61 Y62 Y63 Y64 H65 R66 R69 R70 V71 C77 E78 E79 K80 C84 E87 Q91 I106 Q107 Q115 R122 K127 E130 Q134 V135 R142 Y143 Q189 K190 R167 K171 E172 A173

A174
A175
A176

- Molecule 13: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial

Chain e: 8% 81% 18%

P51 T52 I53 R54 L55 Q56 E57 N70 P71 D72 S73 H74 G75 Y76 D77 K78 D83 F92 S96 S103 Y112 V117 L136 M137 E138 F142 I147 E151 D152 E153 D154

- Molecule 14: NADH dehydrogenase [ubiquinone] 1 subunit C1, mitochondrial

Chain f: 27% 94% 6%

K28 F29 Y30 I31 R32 E33 P34 R36 G37 S38 P39 L42 T47 E76

- Molecule 15: NADH dehydrogenase [ubiquinone] 1 subunit C2

Chain g: 79% 20%

M1 T2 M3 M4 S5 G6 R7 R10 L13 Q14 F15 E19 L33 D47 R52 V55 V56 S57 A58 L65 Y66 Y85 D89 M92 V96 E101 E105 K106 D107 K108 K109 T110 E113 I114 K122

- Molecule 16: NADH dehydrogenase [ubiquinone] iron-sulfur protein 5

Chain h: 10% 85% 15%



P2 M18 T19 I20 Q21 E24 K28 G31 R32 C33 F36 E37 H45 G49 D62 Q70 R81 D84 K85 L86 I87 K88 E89 G90 K91 Y92 L99 E102 D103 P106

- Molecule 17: NADH-ubiquinone oxidoreductase chain 2


Chain i: 74% 24%




- Molecule 18: NADH-ubiquinone oxidoreductase chain 3




- Molecule 19: NADH-ubiquinone oxidoreductase chain 4L




- Molecule 20: NADH-ubiquinone oxidoreductase chain 5





- Molecule 21: NADH-ubiquinone oxidoreductase chain 6







- Molecule 22: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1




- Molecule 26: NADH-ubiquinone oxidoreductase chain 1




- Molecule 27: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8



- Molecule 28: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 7



- Molecule 29: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	462013	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.076	Depositor
Minimum map value	-0.039	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.003	Depositor
Recommended contour level	0.0123	Depositor
Map size (Å)	257.808, 257.808, 257.808	wwPDB
Map dimensions	480, 480, 480	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.5371, 0.5371, 0.5371	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CDL, PEE, 8Q1, ADP, PLX, UQ

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	Q	0.30	0/380	0.45	0/525
2	S	0.31	0/581	0.57	0/781
3	U	0.26	0/664	0.38	0/912
4	V	0.25	0/1042	0.41	0/1411
5	W	0.31	0/973	0.45	0/1312
6	X	0.29	0/701	0.82	2/948 (0.2%)
7	Y	0.24	0/610	0.45	0/836
8	Z	0.24	0/628	0.50	0/850
9	a	0.33	0/1184	0.50	0/1603
10	b	0.28	0/844	0.60	2/1149 (0.2%)
11	c	0.28	0/1371	0.54	3/1875 (0.2%)
12	d	0.28	0/1494	0.51	2/2015 (0.1%)
13	e	0.29	0/891	0.48	0/1210
14	f	0.23	0/385	0.43	0/522
15	g	0.30	0/1036	0.65	2/1401 (0.1%)
16	h	0.33	0/889	0.59	1/1190 (0.1%)
17	i	0.36	0/2773	0.66	5/3768 (0.1%)
18	j	0.32	0/938	0.52	0/1281
19	k	0.38	0/759	0.80	3/1029 (0.3%)
20	l	0.32	0/4926	0.60	10/6700 (0.1%)
21	m	0.33	0/1304	0.66	5/1773 (0.3%)
22	n	0.24	0/491	0.39	0/663
23	o	0.29	0/1092	0.55	3/1481 (0.2%)
24	p	0.28	0/1590	0.43	0/2155
25	r	0.37	0/3723	0.72	15/5078 (0.3%)
26	s	0.40	0/2581	0.87	17/3529 (0.5%)
27	u	0.30	0/1436	0.53	2/1938 (0.1%)
28	v	0.25	0/1052	0.51	0/1411
29	w	0.27	0/2650	0.47	0/3588
All	All	0.32	0/38988	0.60	72/52934 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
10	b	0	1

There are no bond length outliers.

The worst 5 of 72 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
26	s	10	ILE	N-CA-C	12.06	122.90	110.72
25	r	441	ILE	N-CA-C	10.53	121.36	110.62
25	r	5	ILE	N-CA-C	10.38	121.20	110.72
6	X	71	ALA	CA-C-N	9.81	126.72	119.66
6	X	71	ALA	C-N-CA	9.81	126.72	119.66

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
10	b	121	LYS	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Q	363	0	332	6	0
2	S	566	0	561	6	0
3	U	643	0	642	6	0
4	V	1021	0	1025	13	0
5	W	949	0	935	23	0
6	X	691	0	665	25	0
7	Y	584	0	529	27	0
8	Z	611	0	580	13	0
9	a	1151	0	1164	29	0
10	b	819	0	835	19	0
11	c	1315	0	1208	25	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
12	d	1461	0	1429	40	0
13	e	867	0	817	27	0
14	f	377	0	353	4	0
15	g	1005	0	999	30	0
16	h	867	0	871	16	0
17	i	2710	0	2874	103	0
18	j	914	0	951	21	0
19	k	748	0	799	44	0
20	l	4797	0	4935	180	0
21	m	1273	0	1234	54	0
22	n	479	0	486	8	0
23	o	1062	0	1072	22	0
24	p	1534	0	1470	23	0
25	r	3631	0	3839	120	0
26	s	2508	0	2607	76	0
27	u	1398	0	1374	19	0
28	v	1028	0	982	47	0
29	w	2590	0	2553	49	0
30	U	102	0	164	9	0
30	W	41	0	59	8	0
30	i	47	0	71	11	0
30	j	41	0	59	11	0
30	l	137	0	205	40	0
30	r	51	0	82	16	0
31	V	194	0	294	10	0
31	a	100	0	156	15	0
31	i	87	0	124	4	0
31	l	94	0	141	21	0
31	m	100	0	156	10	0
31	r	199	0	307	53	0
31	s	89	0	125	6	0
31	u	55	0	54	5	0
32	X	35	0	0	1	0
33	a	52	0	88	0	0
33	g	52	0	88	4	0
33	j	104	0	176	9	0
33	r	104	0	176	25	0
34	s	38	0	47	1	0
35	w	27	0	11	6	0
36	Q	5	0	0	0	0
36	S	4	0	0	0	0
36	U	2	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
36	W	1	0	0	0	0
36	a	1	0	0	0	0
36	b	1	0	0	0	0
36	c	3	0	0	0	0
36	d	3	0	0	0	0
36	e	1	0	0	2	0
36	h	3	0	0	1	0
36	i	74	0	0	7	0
36	j	33	0	0	5	0
36	k	35	0	0	11	0
36	l	60	0	0	16	0
36	m	16	0	0	2	0
36	n	2	0	0	0	0
36	p	2	0	0	0	0
36	r	91	0	0	10	0
36	s	92	0	0	11	0
36	w	2	0	0	0	0
All	All	40142	0	40704	1011	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 13.

The worst 5 of 1011 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
25:r:104:LEU:CD2	31:r:505:CDL:H861	1.45	1.43
9:a:97:GLU:OE1	12:d:60:ARG:CD	1.71	1.39
25:r:104:LEU:CD2	31:r:505:CDL:C86	2.02	1.37
7:Y:97:LEU:CD2	28:v:107:ARG:HH11	1.36	1.37
9:a:97:GLU:OE1	12:d:60:ARG:HD2	1.08	1.21

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	42/44 (96%)	40 (95%)	2 (5%)	0	100	100
2	S	68/70 (97%)	64 (94%)	4 (6%)	0	100	100
3	U	81/83 (98%)	79 (98%)	2 (2%)	0	100	100
4	V	138/140 (99%)	135 (98%)	3 (2%)	0	100	100
5	W	111/113 (98%)	108 (97%)	3 (3%)	0	100	100
6	X	86/88 (98%)	81 (94%)	4 (5%)	1 (1%)	10	23
7	Y	65/67 (97%)	62 (95%)	3 (5%)	0	100	100
8	Z	78/80 (98%)	72 (92%)	6 (8%)	0	100	100
9	a	136/138 (99%)	132 (97%)	4 (3%)	0	100	100
10	b	94/126 (75%)	86 (92%)	8 (8%)	0	100	100
11	c	154/156 (99%)	144 (94%)	10 (6%)	0	100	100
12	d	173/175 (99%)	169 (98%)	4 (2%)	0	100	100
13	e	102/104 (98%)	98 (96%)	4 (4%)	0	100	100
14	f	47/49 (96%)	41 (87%)	6 (13%)	0	100	100
15	g	120/122 (98%)	115 (96%)	5 (4%)	0	100	100
16	h	103/105 (98%)	100 (97%)	3 (3%)	0	100	100
17	i	345/347 (99%)	334 (97%)	11 (3%)	0	100	100
18	j	113/115 (98%)	108 (96%)	4 (4%)	1 (1%)	14	30
19	k	96/98 (98%)	95 (99%)	1 (1%)	0	100	100
20	l	604/606 (100%)	578 (96%)	25 (4%)	1 (0%)	43	66
21	m	173/175 (99%)	160 (92%)	13 (8%)	0	100	100
22	n	54/56 (96%)	54 (100%)	0	0	100	100
23	o	126/128 (98%)	120 (95%)	6 (5%)	0	100	100
24	p	176/178 (99%)	166 (94%)	9 (5%)	1 (1%)	21	42
25	r	457/459 (100%)	452 (99%)	4 (1%)	1 (0%)	43	66
26	s	316/318 (99%)	309 (98%)	7 (2%)	0	100	100
27	u	169/171 (99%)	163 (96%)	3 (2%)	3 (2%)	6	14
28	v	122/125 (98%)	115 (94%)	7 (6%)	0	100	100
29	w	318/320 (99%)	300 (94%)	18 (6%)	0	100	100
All	All	4667/4756 (98%)	4480 (96%)	179 (4%)	8 (0%)	44	66

5 of 8 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
6	X	136	GLU
20	l	72	GLN
27	u	168	PHE
18	j	40	GLY
27	u	152	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Q	38/38 (100%)	37 (97%)	1 (3%)	40	68
2	S	57/58 (98%)	57 (100%)	0	100	100
3	U	69/69 (100%)	69 (100%)	0	100	100
4	V	101/101 (100%)	100 (99%)	1 (1%)	68	86
5	W	99/99 (100%)	99 (100%)	0	100	100
6	X	74/81 (91%)	58 (78%)	16 (22%)	1	2
7	Y	62/62 (100%)	62 (100%)	0	100	100
8	Z	55/62 (89%)	55 (100%)	0	100	100
9	a	121/121 (100%)	120 (99%)	1 (1%)	73	88
10	b	90/119 (76%)	90 (100%)	0	100	100
11	c	141/141 (100%)	141 (100%)	0	100	100
12	d	155/155 (100%)	152 (98%)	3 (2%)	50	75
13	e	96/96 (100%)	95 (99%)	1 (1%)	68	86
14	f	35/45 (78%)	35 (100%)	0	100	100
15	g	108/109 (99%)	108 (100%)	0	100	100
16	h	93/93 (100%)	93 (100%)	0	100	100
17	i	311/311 (100%)	309 (99%)	2 (1%)	78	91
18	j	100/100 (100%)	99 (99%)	1 (1%)	68	86
19	k	85/85 (100%)	84 (99%)	1 (1%)	63	83

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
20	l	536/540 (99%)	532 (99%)	4 (1%)	76	89
21	m	126/141 (89%)	125 (99%)	1 (1%)	73	88
22	n	53/53 (100%)	53 (100%)	0	100	100
23	o	113/113 (100%)	113 (100%)	0	100	100
24	p	159/159 (100%)	159 (100%)	0	100	100
25	r	410/410 (100%)	408 (100%)	2 (0%)	81	92
26	s	275/275 (100%)	271 (98%)	4 (2%)	57	80
27	u	153/153 (100%)	153 (100%)	0	100	100
28	v	104/111 (94%)	103 (99%)	1 (1%)	68	86
29	w	283/283 (100%)	282 (100%)	1 (0%)	84	93
All	All	4102/4183 (98%)	4062 (99%)	40 (1%)	65	86

5 of 40 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
20	l	119	LYS
26	s	126	LYS
20	l	197	ASP
25	r	114	GLU
26	s	254	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 103 such sidechains are listed below:

Mol	Chain	Res	Type
20	l	226	GLN
24	p	108	HIS
29	w	235	GLN
20	l	328	HIS
20	l	579	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

28 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
31	CDL	i	401	-	86,86,99	1.18	8 (9%)	92,98,111	0.91	4 (4%)
30	PEE	i	402	-	46,46,50	1.22	6 (13%)	49,51,55	1.03	2 (4%)
32	8Q1	X	201	-	32,34,34	2.27	7 (21%)	39,43,43	1.66	10 (25%)
34	UQ	s	402	-	38,38,63	3.64	10 (26%)	48,49,79	2.77	17 (35%)
33	PLX	j	203	-	51,51,51	1.18	4 (7%)	53,59,59	0.65	1 (1%)
30	PEE	r	501	-	50,50,50	1.18	6 (12%)	53,55,55	0.98	2 (3%)
30	PEE	U	102	-	50,50,50	1.18	6 (12%)	53,55,55	0.96	2 (3%)
33	PLX	a	202	-	51,51,51	1.18	4 (7%)	53,59,59	0.64	1 (1%)
30	PEE	U	101	-	50,50,50	1.18	6 (12%)	53,55,55	0.98	2 (3%)
31	CDL	r	504	-	98,98,99	1.10	9 (9%)	104,110,111	0.95	5 (4%)
33	PLX	r	503	-	51,51,51	0.61	0	53,59,59	0.68	0
31	CDL	V	201	-	93,93,99	1.13	9 (9%)	99,105,111	0.87	4 (4%)
31	CDL	s	401	-	88,88,99	1.11	6 (6%)	94,100,111	1.01	5 (5%)
30	PEE	j	201	-	40,40,50	1.17	5 (12%)	43,45,55	1.02	2 (4%)
33	PLX	j	202	-	51,51,51	1.19	2 (3%)	53,59,59	0.64	2 (3%)
31	CDL	a	201	-	99,99,99	0.92	4 (4%)	105,111,111	1.08	7 (6%)
33	PLX	r	502	-	51,51,51	1.16	4 (7%)	53,59,59	0.69	1 (1%)
33	PLX	g	201	-	51,51,51	1.19	3 (5%)	53,59,59	0.63	1 (1%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
30	PEE	l	704	-	45,45,50	1.25	6 (13%)	48,50,55	1.02	2 (4%)
30	PEE	l	703	-	50,50,50	1.17	6 (12%)	53,55,55	0.97	2 (3%)
35	ADP	w	401	-	28,29,29	3.16	9 (32%)	43,45,45	1.90	9 (20%)
31	CDL	u	201	-	54,54,99	1.25	4 (7%)	60,66,111	1.23	5 (8%)
30	PEE	W	201	-	40,40,50	1.16	5 (12%)	43,45,55	1.04	2 (4%)
31	CDL	V	202	-	99,99,99	1.09	8 (8%)	105,111,111	0.90	4 (3%)
30	PEE	l	701	-	39,39,50	1.34	6 (15%)	42,44,55	1.10	3 (7%)
31	CDL	m	201	-	99,99,99	1.09	7 (7%)	105,111,111	0.92	6 (5%)
31	CDL	r	505	-	99,99,99	1.11	8 (8%)	105,111,111	0.83	4 (3%)
31	CDL	l	702	-	93,93,99	0.94	4 (4%)	99,105,111	1.15	7 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
31	CDL	i	401	-	-	47/97/97/110	-
30	PEE	i	402	-	-	15/50/50/54	-
32	8Q1	X	201	-	-	12/41/41/41	-
34	UQ	s	402	-	-	13/33/57/87	0/1/1/1
33	PLX	j	203	-	-	29/55/55/55	-
30	PEE	r	501	-	-	30/54/54/54	-
30	PEE	U	102	-	-	25/54/54/54	-
33	PLX	a	202	-	-	26/55/55/55	-
30	PEE	U	101	-	-	29/54/54/54	-
31	CDL	r	504	-	-	62/109/109/110	-
33	PLX	r	503	-	-	15/55/55/55	-
31	CDL	V	201	-	-	58/104/104/110	-
31	CDL	s	401	-	-	54/99/99/110	-
30	PEE	j	201	-	-	17/44/44/54	-
33	PLX	j	202	-	-	33/55/55/55	-
31	CDL	a	201	-	-	41/110/110/110	-
33	PLX	r	502	-	-	32/55/55/55	-
33	PLX	g	201	-	-	32/55/55/55	-
30	PEE	l	704	-	-	20/49/49/54	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
30	PEE	l	703	-	-	23/54/54/54	-
35	ADP	w	401	-	-	2/16/32/32	0/3/3/3
31	CDL	u	201	-	-	26/65/65/110	-
30	PEE	W	201	-	-	21/44/44/54	-
31	CDL	V	202	-	-	67/110/110/110	-
30	PEE	l	701	-	-	25/43/43/54	-
31	CDL	m	201	-	-	63/110/110/110	-
31	CDL	r	505	-	-	60/110/110/110	-
31	CDL	l	702	-	-	38/104/104/110	-

The worst 5 of 162 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
34	s	402	UQ	C18-C19	9.99	1.56	1.33
34	s	402	UQ	C13-C14	9.65	1.55	1.33
34	s	402	UQ	C23-C24	9.45	1.54	1.33
34	s	402	UQ	C8-C9	9.32	1.54	1.33
35	w	401	ADP	C3'-C4'	-9.00	1.30	1.53

The worst 5 of 112 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
34	s	402	UQ	C7-C8-C9	-7.61	113.72	126.83
34	s	402	UQ	C22-C23-C24	-6.25	113.32	127.62
34	s	402	UQ	C12-C13-C14	-6.06	113.76	127.62
34	s	402	UQ	C17-C18-C19	-6.02	113.84	127.62
35	w	401	ADP	C5-C4-N3	-5.42	119.26	126.72

There are no chirality outliers.

5 of 915 torsion outliers are listed below:




Mol	Chain	Res	Type	Atoms
30	U	101	PEE	C19-C20-C21-C22
30	U	101	PEE	C11-C10-O2-C2
30	U	101	PEE	C1-O3P-P-O1P
30	U	101	PEE	C1-O3P-P-O4P
30	W	201	PEE	C1-O3P-P-O2P

















There are no ring outliers.


27 monomers are involved in 242 short contacts:




Mol	Chain	Res	Type	Clashes	Symm-Clashes
31	i	401	CDL	4	0
30	i	402	PEE	11	0
32	X	201	8Q1	1	0
34	s	402	UQ	1	0
33	j	203	PLX	4	0
30	r	501	PEE	16	0
30	U	102	PEE	3	0
30	U	101	PEE	6	0
31	r	504	CDL	15	0
33	r	503	PLX	21	0
31	V	201	CDL	3	0
31	s	401	CDL	6	0
30	j	201	PEE	11	0
33	j	202	PLX	5	0
31	a	201	CDL	15	0
33	r	502	PLX	4	0
33	g	201	PLX	4	0
30	l	704	PEE	23	0
30	l	703	PEE	10	0
35	w	401	ADP	6	0
31	u	201	CDL	5	0
30	W	201	PEE	8	0
31	V	202	CDL	7	0
30	l	701	PEE	7	0
31	m	201	CDL	10	0
31	r	505	CDL	38	0
31	l	702	CDL	21	0




The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















Ligand UQ s 402	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PLX j 203	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PEE r 501	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PEE U 102	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>













Ligand PLX r 502	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PLX g 201	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PEE l 704	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>
Ligand PEE l 703	
 <p>Bond lengths</p>	 <p>Bond angles</p>
 <p>Torsions</p>	 <p>Rings</p>









5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-31881. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 240



Y Index: 240




Z Index: 240


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 279



Y Index: 216




Z Index: 255


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0123. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 171 nm³; this corresponds to an approximate mass of 154 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.385 Å⁻¹

8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-31881 and PDB model 7VBL. Per-residue inclusion information can be found in section 3 on page 16.

9.1 Map-model overlay [i](#)


This section was not generated.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0123).





























































9.4 Atom inclusion [i](#)



At the recommended contour level, 86% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0123) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.8540	 0.6380
Q	 0.8150	 0.6250
S	 0.8700	 0.6350
U	 0.7620	 0.6040
V	 0.7450	 0.5990
W	 0.8140	 0.6170
X	 0.8380	 0.6110
Y	 0.7070	 0.5640
Z	 0.6990	 0.5660
a	 0.8790	 0.6550
b	 0.8090	 0.5990
c	 0.8660	 0.6440
d	 0.8240	 0.6230
e	 0.8320	 0.6200
f	 0.6860	 0.5760
g	 0.9010	 0.6560
h	 0.8180	 0.6150
i	 0.9430	 0.6820
j	 0.8040	 0.6370
k	 0.9600	 0.6820
l	 0.9060	 0.6620
m	 0.7860	 0.6100
n	 0.7520	 0.5750
o	 0.8780	 0.6460
p	 0.8710	 0.6360
r	 0.9350	 0.6750
s	 0.9010	 0.6600
u	 0.8020	 0.6090
v	 0.7060	 0.5580
w	 0.8430	 0.6320

