



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 10, 2026 – 07:54 AM UTC

PDB ID : 3VS8 / pdb\_00003vs8  
Title : Crystal structure of type III PKS ArsC  
Authors : Satou, R.; Miyanaga, A.; Ozawa, H.; Funa, N.; Miyazono, K.; Tanokura, M.;  
Ohnishi, Y.; Horinouchi, S.  
Deposited on : 2012-04-23  
Resolution : 1.76 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	NOT EXECUTED
Xtriage (Phenix)	:	2.0
EDS	:	NOT EXECUTED
Buster-report	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.76 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ .

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	410	
1	B	410	
1	C	410	
1	D	410	
1	E	410	
1	F	410	
1	G	410	
1	H	410	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 27375 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Type III polyketide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	404	Total	C	N	O	S	0	0	0
			3123	1984	542	582	15			
1	B	399	Total	C	N	O	S	0	0	0
			3084	1961	537	571	15			
1	C	402	Total	C	N	O	S	0	0	0
			3108	1973	540	580	15			
1	D	401	Total	C	N	O	S	0	0	0
			3100	1969	539	577	15			
1	E	401	Total	C	N	O	S	0	0	0
			3107	1976	539	577	15			
1	F	404	Total	C	N	O	S	0	0	0
			3123	1984	542	582	15			
1	G	398	Total	C	N	O	S	0	0	0
			3078	1958	536	569	15			
1	H	384	Total	C	N	O	S	0	0	0
			2968	1891	518	544	15			

- Molecule 2 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Na	0	0
			1	1		
2	B	1	Total	Na	0	0
			1	1		
2	C	1	Total	Na	0	0
			1	1		
2	D	1	Total	Na	0	0
			1	1		
2	E	1	Total	Na	0	0
			1	1		
2	F	1	Total	Na	0	0
			1	1		

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	G	1	Total 1	Na 1	0	0
2	H	1	Total 1	Na 1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	437	Total 437	O 437	0	0
3	B	258	Total 258	O 258	0	0
3	C	361	Total 361	O 361	0	0
3	D	356	Total 356	O 356	0	0
3	E	417	Total 417	O 417	0	0
3	F	330	Total 330	O 330	0	0
3	G	283	Total 283	O 283	0	0
3	H	234	Total 234	O 234	0	0

### 3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


#### • Molecule 1: Type III polyketide synthase




#### • Molecule 1: Type III polyketide synthase




#### • Molecule 1: Type III polyketide synthase






- Molecule 1: Type III polyketide synthase

Chain D: 78% 17% ..




- Molecule 1: Type III polyketide synthase

Chain E: 80% 17% ..



- Molecule 1: Type III polyketide synthase


Chain F: 78% 18% ..



- Molecule 1: Type III polyketide synthase


Chain G: 76% 18% ..





• Molecule 1: Type III polyketide synthase

Chain H: 69% 20% 6%



## 4 Data and refinement statistics

EDS was not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	102.32Å 143.39Å 129.62Å 90.00° 110.33° 90.00°	Depositor
Resolution (Å)	37.57 – 1.76	Depositor
% Data completeness (in resolution range)	97.4 (37.57-1.76)	Depositor
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.75 (at 1.76Å)	Xtriage
Refinement program	REFMAC 5.5.0066	Depositor
R, $R_{free}$	0.183 , 0.230	Depositor
Wilson B-factor (Å <sup>2</sup> )	22.9	Xtriage
Anisotropy	0.179	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.023 for h,-k,-h-l	Xtriage
Total number of atoms	27375	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 72.42 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.2090e-06. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality ⓘ

### 5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	1.66	35/3194 (1.1%)	1.35	10/4331 (0.2%)
1	B	1.44	19/3152 (0.6%)	1.34	22/4270 (0.5%)
1	C	1.54	26/3177 (0.8%)	1.30	14/4307 (0.3%)
1	D	1.48	19/3169 (0.6%)	1.29	17/4296 (0.4%)
1	E	1.55	18/3177 (0.6%)	1.33	11/4307 (0.3%)
1	F	1.49	23/3194 (0.7%)	1.35	17/4331 (0.4%)
1	G	1.31	7/3148 (0.2%)	1.25	10/4267 (0.2%)
1	H	1.32	12/3034 (0.4%)	1.32	15/4110 (0.4%)
All	All	1.48	159/25245 (0.6%)	1.32	116/34219 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	F	0	1
1	G	0	2
All	All	0	3

The worst 5 of 159 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	299	VAL	CA-CB	-16.56	1.45	1.54
1	A	373	ILE	CA-CB	14.55	1.62	1.54
1	B	195	MET	C-O	10.43	1.36	1.24
1	E	186	CYS	CA-CB	10.26	1.70	1.53
1	C	222	MET	SD-CE	-10.01	1.54	1.79

The worst 5 of 116 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	276	PRO	CA-C-N	-17.58	99.47	123.03
1	H	276	PRO	C-N-CA	-17.58	99.47	123.03
1	F	103	ASP	N-CA-C	10.19	126.01	113.17
1	E	186	CYS	CA-CB-SG	-9.58	92.37	114.40
1	E	298	LEU	CA-C-N	-9.56	114.22	120.24

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	F	103	ASP	Peptide
1	G	100	PRO	Peptide
1	G	92	ASP	Peptide

## 5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3123	0	3112	30	0
1	B	3084	0	3083	88	0
1	C	3108	0	3099	72	0
1	D	3100	0	3095	57	0
1	E	3107	0	3101	46	0
1	F	3123	0	3112	66	0
1	G	3078	0	3074	80	0
1	H	2968	0	2986	104	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
2	E	1	0	0	0	0
2	F	1	0	0	0	0
2	G	1	0	0	0	0
2	H	1	0	0	0	0
3	A	437	0	0	8	1
3	B	258	0	0	13	0
3	C	361	0	0	8	1
3	D	356	0	0	14	2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	E	417	0	0	10	0
3	F	330	0	0	13	0
3	G	283	0	0	11	0
3	H	234	0	0	7	0
All	All	27375	0	24662	531	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:89:ARG:CB	1:C:96:ASP:HB3	1.55	1.35
1:C:89:ARG:HB2	1:C:96:ASP:CB	1.62	1.29
1:D:122:LYS:HD3	3:D:872:HOH:O	1.32	1.26
1:C:39:ARG:NH2	1:C:94:ASN:ND2	1.86	1.22
1:H:104:GLN:HE21	1:H:104:GLN:CA	1.51	1.22

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:864:HOH:O	3:D:864:HOH:O[1_655]	2.04	0.16
3:A:998:HOH:O	3:D:946:HOH:O[2_646]	2.16	0.04

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	402/410 (98%)	391 (97%)	11 (3%)	0	100	100
1	B	395/410 (96%)	381 (96%)	14 (4%)	0	100	100

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	C	398/410 (97%)	386 (97%)	11 (3%)	1 (0%)	36	21
1	D	397/410 (97%)	385 (97%)	12 (3%)	0	100	100
1	E	397/410 (97%)	386 (97%)	11 (3%)	0	100	100
1	F	402/410 (98%)	389 (97%)	12 (3%)	1 (0%)	43	27
1	G	394/410 (96%)	376 (95%)	16 (4%)	2 (0%)	24	11
1	H	378/410 (92%)	360 (95%)	18 (5%)	0	100	100
All	All	3163/3280 (96%)	3054 (97%)	105 (3%)	4 (0%)	48	32

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	F	104	GLN
1	G	92	ASP
1	C	102	PRO
1	G	102	PRO

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	333/338 (98%)	327 (98%)	6 (2%)	51	33
1	B	328/338 (97%)	317 (97%)	11 (3%)	32	13
1	C	332/338 (98%)	316 (95%)	16 (5%)	23	6
1	D	331/338 (98%)	321 (97%)	10 (3%)	36	16
1	E	332/338 (98%)	323 (97%)	9 (3%)	39	19
1	F	333/338 (98%)	319 (96%)	14 (4%)	26	8
1	G	328/338 (97%)	316 (96%)	12 (4%)	30	11
1	H	317/338 (94%)	298 (94%)	19 (6%)	17	3
All	All	2634/2704 (97%)	2537 (96%)	97 (4%)	30	11

5 of 97 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	F	150	LEU
1	G	226	ASN
1	F	231	SER
1	F	373	ILE
1	G	346	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
1	E	343	HIS
1	G	204	GLN
1	F	78	GLN
1	G	78	GLN
1	H	41	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

Mogul was not executed - this section is therefore empty.

## 5.5 Carbohydrates [i](#)

Mogul was not executed - this section is therefore empty.

## 5.6 Ligand geometry [i](#)

Mogul was not executed - this section is therefore empty.

## 5.7 Other polymers [i](#)

Mogul was not executed - this section is therefore empty.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

### 6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

### 6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

### 6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.