



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 12, 2026 – 05:00 PM UTC

PDB ID : 5WER / pdb\_00005wer  
Title : Crystal Structure of TAPBPR and H2-Dd complex  
Authors : Jiang, J.S.; Natarajan, K.; Boyd, L.F.; Margulies, D.H.  
Deposited on : 2017-07-10  
Resolution : 3.41 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.41 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	1210 (3.48-3.36)
Clashscore	190562	1234 (3.48-3.36)
Ramachandran outliers	187476	1222 (3.48-3.36)
Sidechain outliers	187428	1222 (3.48-3.36)
RSRZ outliers	180081	1210 (3.48-3.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	277	 69% 27% .
1	D	277	 72% 23% .
1	G	277	 74% 22% ..
1	J	277	 73% 24% .
2	B	100	 74% 24% .

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	E	100	
2	H	100	
2	K	100	
3	C	394	
3	F	394	
3	I	394	
3	L	394	

## 2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 20155 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called H-2 class I histocompatibility antigen, D-D alpha chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	265	Total	C	N	O	S	256	0	0
			2086	1310	374	393	9			
1	D	265	Total	C	N	O	S	82	0	0
			1973	1231	354	379	9			
1	G	269	Total	C	N	O	S	132	0	0
			2004	1252	364	379	9			
1	J	270	Total	C	N	O	S	173	0	0
			2047	1283	364	390	10			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	1	MET	-	initiating methionine	UNP P01900
A	73	CYS	SER	engineered mutation	UNP P01900
D	1	MET	-	initiating methionine	UNP P01900
D	73	CYS	SER	engineered mutation	UNP P01900
G	1	MET	-	initiating methionine	UNP P01900
G	73	CYS	SER	engineered mutation	UNP P01900
J	1	MET	-	initiating methionine	UNP P01900
J	73	CYS	SER	engineered mutation	UNP P01900

- Molecule 2 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	98	Total	C	N	O	S	32	0	0
			774	490	131	151	2			
2	E	98	Total	C	N	O	S	20	0	0
			771	488	128	153	2			
2	H	98	Total	C	N	O	S	14	0	0
			781	497	128	154	2			
2	K	98	Total	C	N	O	S	15	0	0
			785	498	133	152	2			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	initiating methionine	UNP P61769
E	0	MET	-	initiating methionine	UNP P61769
H	0	MET	-	initiating methionine	UNP P61769
K	0	MET	-	initiating methionine	UNP P61769

- Molecule 3 is a protein called TAP binding protein related.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	317	Total	C	N	O	S	120	0	0
			2362	1490	401	460	11			
3	F	315	Total	C	N	O	S	40	0	0
			2182	1369	371	433	9			
3	I	301	Total	C	N	O	S	56	0	0
			2092	1307	356	418	11			
3	L	312	Total	C	N	O	S	36	0	0
			2243	1407	383	443	10			

- Molecule 4 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C<sub>2</sub>H<sub>6</sub>O<sub>2</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		
4	E	1	Total	C	O	0	0
			4	2	2		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	C	1	Total	C	O	0	0
			6	3	3		
5	F	1	Total	C	O	0	0
			6	3	3		
5	L	1	Total	C	O	0	0
			6	3	3		

- Molecule 6 is PENTAETHYLENE GLYCOL (CCD ID: 1PE) (formula:  $C_{10}H_{22}O_6$ ).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	J	1	Total	C	O	0	0
			16	10	6		

- Molecule 7 is CITRIC ACID (CCD ID: CIT) (formula:  $C_6H_8O_7$ ).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
7	L	1	Total	C	O	0	0
			13	6	7		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: H-2 class I histocompatibility antigen, D-D alpha chain



- Molecule 1: H-2 class I histocompatibility antigen, D-D alpha chain




- Molecule 1: H-2 class I histocompatibility antigen, D-D alpha chain



- Molecule 1: H-2 class I histocompatibility antigen, D-D alpha chain




Chain J:  73% 24% .



- Molecule 2: Beta-2-microglobulin

Chain B:  74% 24% .




- Molecule 2: Beta-2-microglobulin

Chain E:  64% 34% .




- Molecule 2: Beta-2-microglobulin

Chain H:  77% 21% .




- Molecule 2: Beta-2-microglobulin


Chain K:  62% 36% .



- Molecule 3: TAP binding protein related


Chain C:  51% 28% 20% .






• Molecule 3: TAP binding protein related


Chain F: 52% 25% 20%




• Molecule 3: TAP binding protein related

Chain I: 51% 25% 24%





● Molecule 3: TAP binding protein related



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 31	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	169.05Å 169.05Å 139.41Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	48.80 – 3.41 48.80 – 3.41	Depositor EDS
% Data completeness (in resolution range)	97.2 (48.80-3.41) 97.2 (48.80-3.41)	Depositor EDS
$R_{merge}$	0.42	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.91 (at 3.40Å)	Xtriage
Refinement program	PHENIX (1.10.1_2155)	Depositor
R, $R_{free}$	0.239 , 0.269 0.239 , 0.273	Depositor DCC
$R_{free}$ test set	2948 reflections (4.89%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	83.1	Xtriage
Anisotropy	0.304	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 604.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.379 for -h,-k,l 0.046 for h,-h-k,-l 0.045 for -k,-h,-l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	20155	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	76.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.40% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, 1PE, CIT, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.16	0/2142	0.43	0/2916
1	D	0.13	0/2023	0.34	0/2762
1	G	0.13	0/2054	0.35	0/2802
1	J	0.12	0/2100	0.35	0/2866
2	B	0.12	0/795	0.36	0/1083
2	E	0.16	0/791	0.38	0/1079
2	H	0.12	0/803	0.34	0/1095
2	K	0.12	0/807	0.33	0/1100
3	C	0.20	0/2397	0.45	0/3263
3	F	0.25	0/2212	0.53	0/3026
3	I	0.19	0/2120	0.46	0/2897
3	L	0.21	0/2272	0.54	0/3098
All	All	0.17	0/20516	0.43	0/27987

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2086	0	1873	57	0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	D	1973	0	1695	43	0
1	G	2004	0	1732	41	0
1	J	2047	0	1794	53	0
2	B	774	0	693	17	0
2	E	771	0	698	31	0
2	H	781	0	710	15	0
2	K	785	0	717	32	0
3	C	2362	0	2337	74	0
3	F	2182	0	2007	72	0
3	I	2092	0	1918	70	0
3	L	2243	0	2167	78	0
4	A	4	0	6	0	0
4	E	4	0	6	0	0
5	C	6	0	7	0	0
5	F	6	0	8	0	0
5	L	6	0	7	1	0
6	J	16	0	22	0	0
7	L	13	0	5	0	0
All	All	20155	0	18402	531	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

The worst 5 of 531 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:J:111:ARG:NH1	3:L:213:ARG:HH12	1.71	0.88
1:J:119:ASP:OD2	2:K:1:ILE:N	2.08	0.85
3:I:149:LEU:HD13	3:I:156:LEU:HB3	1.59	0.84
3:F:13:ASP:N	3:F:44:LYS:O	2.15	0.79
3:C:14:VAL:HB	3:C:43:LEU:HB2	1.65	0.79

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/277 (94%)	234 (90%)	22 (8%)	3 (1%)	10	35
1	D	259/277 (94%)	238 (92%)	16 (6%)	5 (2%)	6	26
1	G	263/277 (95%)	233 (89%)	24 (9%)	6 (2%)	5	23
1	J	266/277 (96%)	241 (91%)	22 (8%)	3 (1%)	11	37
2	B	96/100 (96%)	86 (90%)	9 (9%)	1 (1%)	12	39
2	E	96/100 (96%)	85 (88%)	11 (12%)	0	100	100
2	H	96/100 (96%)	88 (92%)	8 (8%)	0	100	100
2	K	96/100 (96%)	86 (90%)	10 (10%)	0	100	100
3	C	299/394 (76%)	254 (85%)	36 (12%)	9 (3%)	3	18
3	F	297/394 (75%)	247 (83%)	33 (11%)	17 (6%)	1	9
3	I	283/394 (72%)	216 (76%)	58 (20%)	9 (3%)	3	18
3	L	294/394 (75%)	234 (80%)	44 (15%)	16 (5%)	1	10
All	All	2604/3084 (84%)	2242 (86%)	293 (11%)	69 (3%)	4	21

5 of 69 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	184	PRO
3	C	70	PRO
3	C	307	PRO
3	F	157	PRO
3	F	230	LYS

### 5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	197/232 (85%)	197 (100%)	0	100	100
1	D	174/232 (75%)	174 (100%)	0	100	100

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	176/232 (76%)	176 (100%)	0	100	100
1	J	184/232 (79%)	184 (100%)	0	100	100
2	B	81/95 (85%)	80 (99%)	1 (1%)	63	72
2	E	83/95 (87%)	83 (100%)	0	100	100
2	H	85/95 (90%)	85 (100%)	0	100	100
2	K	85/95 (90%)	85 (100%)	0	100	100
3	C	259/325 (80%)	259 (100%)	0	100	100
3	F	212/325 (65%)	212 (100%)	0	100	100
3	I	206/325 (63%)	206 (100%)	0	100	100
3	L	235/325 (72%)	235 (100%)	0	100	100
All	All	1977/2608 (76%)	1976 (100%)	1 (0%)	88	88

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	38	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	H	42	ASN
3	I	45	GLN
3	L	263	GLN
3	L	155	ASN
3	L	177	GLN

### 5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

7 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
5	GOL	L	402	-	5,5,5	0.37	0	5,5,5	0.31	0
5	GOL	F	401	-	5,5,5	0.40	0	5,5,5	0.23	0
5	GOL	C	401	-	5,5,5	0.39	0	5,5,5	0.32	0
4	EDO	A	301	-	3,3,3	0.43	0	2,2,2	0.37	0
6	1PE	J	301	-	15,15,15	0.55	0	14,14,14	0.22	0
4	EDO	E	101	-	3,3,3	0.43	0	2,2,2	0.37	0
7	CIT	L	401	-	12,12,12	1.08	0	17,17,17	1.58	3 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	GOL	L	402	-	-	2/4/4/4	-
5	GOL	F	401	-	-	2/4/4/4	-
5	GOL	C	401	-	-	2/4/4/4	-
4	EDO	A	301	-	-	0/1/1/1	-
6	1PE	J	301	-	-	7/13/13/13	-
4	EDO	E	101	-	-	0/1/1/1	-
7	CIT	L	401	-	-	12/16/16/16	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
7	L	401	CIT	O6-C6-C3	3.83	120.49	113.14
7	L	401	CIT	O2-C1-C2	2.30	121.63	114.35
7	L	401	CIT	O4-C5-C4	2.13	121.08	114.35

There are no chirality outliers.

5 of 25 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	L	402	GOL	O1-C1-C2-C3
7	L	401	CIT	O7-C3-C4-C5
7	L	401	CIT	C6-C3-C4-C5
7	L	401	CIT	O7-C3-C6-O5
7	L	401	CIT	O7-C3-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	L	402	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	265/277 (95%)	-1.36	0 100 100	7, 76, 107, 121	78 (29%)
1	D	265/277 (95%)	-1.39	0 100 100	24, 77, 105, 120	27 (10%)
1	G	268/277 (96%)	-1.36	0 100 100	24, 82, 113, 127	36 (13%)
1	J	260/277 (93%)	-1.33	0 100 100	10, 84, 113, 133	34 (13%)
2	B	98/100 (98%)	-1.44	0 100 100	21, 70, 105, 117	6 (6%)
2	E	98/100 (98%)	-1.34	0 100 100	44, 74, 108, 118	8 (8%)
2	H	98/100 (98%)	-1.42	0 100 100	44, 73, 113, 129	4 (4%)
2	K	98/100 (98%)	-1.36	0 100 100	42, 79, 98, 115	3 (3%)
3	C	317/394 (80%)	-1.39	0 100 100	23, 64, 102, 128	32 (10%)
3	F	315/394 (79%)	-1.40	0 100 100	27, 70, 101, 129	14 (4%)
3	I	299/394 (75%)	-1.39	0 100 100	18, 66, 94, 113	13 (4%)
3	L	312/394 (79%)	-1.46	0 100 100	6, 60, 95, 124	13 (4%)
All	All	2693/3084 (87%)	-1.39	0 100 100	6, 71, 107, 133	268 (9%)

There are no RSRZ outliers to report.

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
5	GOL	L	402	6/6	0.97	0.04	66,88,94,101	0
4	EDO	E	101	4/4	0.98	0.06	57,61,66,66	0
4	EDO	A	301	4/4	0.98	0.03	55,65,66,69	0
6	1PE	J	301	16/16	0.98	0.04	50,61,89,91	0
5	GOL	C	401	6/6	0.99	0.02	30,47,56,74	1
5	GOL	F	401	6/6	0.99	0.07	24,39,43,46	0
7	CIT	L	401	13/13	0.99	0.02	20,41,59,59	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.